



**COTSWOLD**  
Grass Seeds  
— DIRECT —

**2021**

# SEED CATALOGUE

48<sup>TH</sup> EDITION

[cotswoldseeds.com](http://cotswoldseeds.com)

# Experts in their field

- The right advice
- Bespoke mixtures



Lizzie

Sam

James

Abbi

Call 01608 652552

Visit [cotswoldseeds.com](http://cotswoldseeds.com)

## Reading the mixtures

■ 2.00 kg certified TODDINGTON perennial ryegrass.  
The green block indicates that this is a grass.



This bar would indicate a mixture of 50% grass and 50% legume content - based on weight.



You will see a key on every page where there is a mixture, showing which colour represents which 'type' of plant.

## Payment

Account customers receive 28 days credit.  
Payment can also be made by card, cheque, BACS and PayPal.

Payable to: Cotswold Seeds Ltd  
Barclays Bank  
Acc. no. 80278661  
Sort code 20-20-15

IBAN: GB92 BARC 2020 15802786 61  
SWIFTBIC: BARCGB22

T&Cs apply - available on website/invoices.

Customers can now order online with clarification of farmers group membership.

## Free delivery on orders over £75

Orders placed by 2pm on the phone or online can normally be delivered next day to most parts of the UK (not weekends & stock permitting). We try and use our own vehicles for delivery where possible, and use third party carriers for all other areas. We always endeavour to mix your seed and dispatch on the day of ordering prior to 2pm.

We also deliver to the outlying areas of the UK and Europe; any additional cost will be advised at the time of ordering. Customers outside the UK, Northern Ireland & Eire should order by phone, rather than online.

N.B. In the event of shortages we reserve the right to use alternative varieties in our mixes without notice. Please check website for latest updates.

© Copyright: Cotswold Seeds Ltd  
ISBN: 978-1-8380163-1-9







# Hand Picked & Mixed

Next day delivery

Since 1974

It's an understatement to say that 2020 was a challenging year for us all, but we are looking forward to a much better season ahead. I'm pleased to say that seed supply has recovered from the difficult growing seasons in 2018 and 2019, so this will help us get off to a good start, with a great selection of species and varieties secured for the coming year.

Before we move on we would like to thank you, our loyal customers, for your patience in these very challenging circumstances, but also a big thank you to our staff and suppliers for pulling out all the stops for us in 2020. Over the decades, Cotswold Seeds has built its reputation on first class customer care. It's hugely important to us, from offering helpful, friendly and impartial technical advice, to next day delivery. We've worked very hard to maintain this level of service and the attention to detail that makes life run that little bit more smoothly. Thankfully, farms were able to continue operating during the pandemic, and we were able to do the same, implementing many measures to keep the mixers rolling and the seed flowing safely. Throughout the busy periods our warehouse team were working 24/7 to overcome bottlenecks and satisfy demand. A huge thanks to the finest seed mixing team in the land!

Although some of the customer days at FarmED were postponed in 2020, we have rescheduled these for spring and summer 2021. Our MD, Ian Wilkinson, will be thrilled to welcome you on a farm walk followed by some delicious and nutritious sustainable food from the amazing new FarmED kitchen.

We wish you every success in 2021, hoping for great growing weather and good health. Please rest assured that whatever the next 12 months has in store, we will be here for you, ready to advise, mix and deliver.

Thank you for your continued support.

*Paul Totterdell*  
General Manager

## INDEX

OVER-SEEDING	08
SILAGE & HAY	10
GRAZING & FORAGE	18
ROOT CROPS	30
EQUINE	34
SOIL	36
ENVIRONMENTAL	41
WILD FLOWERS	46
GAME	50
LAWN & LANDSCAPE	54



# Grasses

Grassland is the single most important source of forage for British farmers.

Our climate is ideal for grass growth, making grazed grass easily the cheapest source of forage for livestock. In order to capitalise on this great natural resource, extensive research over many decades has improved UK grassland productivity and its on-farm utilisation dramatically.

However, of the 50 or so different types of grass found in the UK, only a handful are cultivated on any scale, with the most important outlined here.

Ryegrass, which comes in many different forms, is the most widely sown of all grasses. Ryegrasses have high sugars and respond to nitrogen fertiliser better than any other grass species. These two qualities have made it the most popular grass for silage over the last sixty years (since the Plough Up policy of WW2 and the advent of cheap nitrogen fertiliser).

Increased demand resulted in the development of new varieties led by Sir George Stapledon at the Aberystwyth Plant Breeding Station. Other plant breeders across Europe followed suit and, as a result, we now have a comprehensive range of varieties to select from.

## How Long Do Ryegrass Leys Last?

Perennial ryegrass based leys last between three and five years reliably. On good soils they can last longer, but all eventually deteriorate as unsown species such as meadowgrasses and bents increase to make up more of the sward.

There are differences within ryegrass species and between individual varieties. Generally, late heading perennial ryegrasses such as Toddington are very persistent with good ground cover. Earlier heading ryegrasses such as Kirial, a hybrid type, offer early season growth but do not persist as well.

In all circumstances, ryegrass leys should be considered temporary and should not be routinely extended beyond the duration recommended for each mixture. Over-seeding is a good way to prolong their life (see page 8).

### 1 Perennial Ryegrass (*Lolium perenne*)

This is the most persistent type of ryegrass and by far the most widely sown. It yields around 13t DM per hectare which is lower than Italian ryegrass. However, it is more flexible in use because it can be grazed or cut and made into silage, haylage or hay. There are many varieties to choose from, some are very leafy with little stem and are excellent for grazing, others have much earlier, upright growth which make them well suited to silage making. Most perennial ryegrasses last around five years or more.

### 2 Westerwolds Ryegrass (*Lolium westerwoldicum*)

Westerwolds is the highest yielding ryegrass with similar forage quality to the well known Italian ryegrass. Westerwolds is capable of extremely fast growth, is generous in response to nitrogen fertiliser and is grown largely for silage production. It is an annual, surviving for one season only. It may be sown in the autumn for production the following spring and summer, or planted in the spring for summer cropping. When sown in the spring it is ready for cutting after only 12 weeks and further growth will follow where soil moisture is plentiful.

### 3 Italian Ryegrass (*Lolium multiflorum*)

This is a short lived grass lasting for two years. It is very high yielding and reliably provides up to 18t DM per hectare on soils that suit it. (All ryegrasses yield less on light soils, especially in low rainfall areas.) It has a very open growth habit with fewer tillers than other grasses and is therefore better suited to cutting than grazing. Modern varieties offer high yields and good disease resistance.

### 4 Hybrid Ryegrass (*Lolium x boucheanum*)

This form of ryegrass is perhaps one of the best grasses available to the intensive farmer. The hybrid is a cross between the Italian and perennial forms of ryegrass and shares characteristics of both. The dominant parent determines how the variety performs in the field. Most hybrid varieties have the Italian gene dominant and the best cultivars provide the same or similar high yields as Italian ryegrass. But, as they also contain some of the persistent genes

of the perennial ryegrass parent, they last longer. The genes of the perennial ryegrass parent produce a plant with more tillers and more leaf which gives increased ground cover, making it better for grazing.

### Tetraploid ryegrass

Modern plant breeding has produced tetraploid ryegrass varieties. These are available in Italian, hybrid and perennial form. With double the number of chromosomes of the standard diploid varieties their characteristics differ. Tetraploid ryegrasses are highly palatable which leads to higher voluntary intakes, of great value in seed mixtures. However, they also tiller less than diploids which means that they do not cover so much ground, leaving more soil showing. They are also less persistent. For these reasons, tetraploids should be used at low levels in long term grazing leys but can be used at higher levels in silage leys.

### 5 Cocksfoot (*Dactylis glomerata*)

Of all the grasses, cocksfoot has the deepest roots and, when grown on dry or free-draining soil, offers continued growth in dry weather while adding plenty of organic matter to hungry, thin soils. Cocksfoot provides 'early bite' in spring and quick recovery after grazing or cutting. It is very good for up to four years provided it is grazed hard as it will then remain leafy. However, cocksfoot is not a grass to choose for long term pasture as it tends to become clumpy, coarse and unpalatable.

### 6 Timothy (*Phleum pratense*)

Possibly the most important long term agricultural grass, timothy is commonly found in pasture throughout the UK. It will grow abundantly on heavy ground and, although it only has a shallow root structure, persists well on lighter land in dry years. It is very persistent and disease free. The forage it produces is acceptable to most stock and it can be made into silage and hay or grazed. Another form of timothy, smaller catstail (*Phleum bertolonii*), is shorter, less dominant and lower yielding but is a useful component of mixtures for environmental purposes.



## 7 Meadow Fescue (*Festuca pratensis*)

A long duration grass that is often sown with timothy to provide hay or grazing. For longer term leys it is an alternative to perennial ryegrass, especially in upland areas. It will grow on nearly all soils ranging from light, brashy types to stiff clays. It has the same growth habit as perennial ryegrass and, although more persistent and drought tolerant, is slower to establish.

### Festulolium

A recent development in plant breeding has produced this natural hybridisation of ryegrass and fescue, combining the stress resistant genes of fescue with the bulky yield of ryegrass, improving drought resistance with high yield.

## 8 Common Bent (*Agrostis capillaris*)

This delicately flowered grass is included in the majority of agri-environmental mixes. As it has a tiny seed it is added to mixes at low levels. It is a creeping grass and, although of little agricultural value, is very common in old grasslands. It is adaptable to most soils and is drought tolerant.

## 9 Creeping Red Fescue (*Festuca rubra rubra*)

This common grass has creeping roots which enable it to remain green in dry times and give pasture a good bottom. Sometimes this can also be a disadvantage as it stifles some of the more delicate species and should therefore be used with caution. An alternative fescue, such as sheeps, red or slender creeping red will allow the

development of finer species. However, creeping red fescue is an inexpensive seed and can be included in simple mixtures, particularly those for low grade amenity use.

## 10 Meadow Foxtail (*Alopecurus pratensis*)

A tufted perennial which is widespread throughout the British Isles. It is commonly found in low-lying areas, particularly river meadows. Nutritious and palatable to stock, it is one of the first grasses to flower in the spring. When making hay, it makes a useful contribution to yields.

## 11 Red Fescue (*Festuca rubra commutata*)

Also known as chewings fescue, this is a fine leaved, tufted grass. It is distinguished from creeping red fescue by an absence of creeping rhizomes. It tolerates drought well and is common on well-drained, gravelly, chalky and sandy soils in the south. It forms a dense turf and is one of the main species used with bent to form lawns.

## 12 Sheeps Fescue (*Festuca ovina*)

The finest leaved and least aggressive fescue which allows other delicate species room to establish. It only grows to 15 – 25cm, is very hardy and can be found in all areas of the UK. Although it provides only low levels of production, the forage it produces is of reasonable quality. It will grow on most soils and tolerates low fertility situations.

## 13 Crested Dogstail (*Cynosurus cristatus*)

Traditionally a grazing grass, this compact, tufted perennial is found in abundance in sheep pastures. It is not aggressive and grows well late into the season when other grasses are giving up. It grows in most areas, even on clay soils, but is found naturally in dry areas. It has good winter greenness but is inclined to produce wiry stems if not cut or grazed.

## 14 Smooth Stalked Meadowgrass (*Poa pratensis*)


This perennial has creeping rhizomes and is very drought resistant. It is common throughout the UK, particularly on lighter soils. It should not be sown late in the autumn as it is slow to establish. Shallow sowing is also essential as the seed needs light to germinate. Early to grow in the spring, once topped or cut it tends not to re-flower so regrowth is leafy.

## 15 Sweet Vernal Grass (*Anthoxanthum odoratum*)

An early flowering grass, strongly scented with coumarin, often found in old pastures and meadows and sometimes included in seed mixtures to give scent to hay. It has a high proportion of stem to leaf and so is unpalatable to stock. It is an attractive grass but seed is expensive so is usually included at a low levels in seed mixtures.

## 16 Tall Fescue (*Festuca arundinacea*)

The largest fescue which forms sizable, dense tussocks. It can grow to six feet tall, particularly on damp or wet soils. On light soils it is drought resistant but it is less palatable than meadow fescue and so is less attractive to farmers for forage production.





# Legumes

Legumes, grown with grass or on their own, play an important role in providing highly nutritious forage and free nitrogen.

All legumes share the ability to collect nitrogen from the air and make it available in the soil for plant growth.

Legume-rich forage is therefore low cost as it requires little or no nitrogen fertiliser. Legumes are also high in protein and, because they are particularly relished by livestock, improve animal performance.

There are twelve legumes commonly used including the true clovers, the medics, sainfoin, birdsfoot trefoil and vetches.

## True Clovers

### 1 White Clover (*Trifolium repens*)

White clover is probably one of the most valuable plants in existence and is the most popular forage legume. It differs from other clovers in having a stolon (or stem) that runs along the ground. This produces edible leaves and flower heads at low levels, making it ideal for grazing. It is long lasting and drought resistant and grows on nearly all soils. White clover has received more research funding than any other legume and so is well understood. In common with most fodder legumes, it is best grown with grasses which increase total forage yield and produce a flexible sward which can be cut or grazed.

#### Increase livestock productivity

White clover has a high protein content at around 20-25%. Perennial ryegrass contains about 16%. Combining these two together in the field increases the overall protein content of forage by 2-3% to around 20%.

The extra protein available from clover leys has a direct impact on live weight gains. At the same time, grazing animals consume more as they find clover very palatable. This all results in animals fattening faster compared to those on non-clover leys.

#### A leaf size for every purpose

There is a large range of white clovers available, classified by leaf size, with the tolerance for close grazing increasing as leaf size decreases. Medium-leaved varieties, such as AberSwan and AberDai, are good for grazing, silage or hay. Large-leaved strains, such as Alice, give slightly higher yields but are less persistent when grazed and are therefore for cutting only.

### 2 Red Clover (*Trifolium pratense*)

Red clover produces a third more yield than white clover but is less persistent, only lasting for between two and four years. It is normally used to produce silage, although it can be grazed occasionally.

It is an erect and dominant plant that is best sown with aggressive ryegrasses. However, it may be included in more complex seed mixes but its inclusion rate must be low to counter its aggression. It grows on nearly all soils except acidic ones where alsike clover should be used.

#### Oestrogen and livestock fertility

Red clover contains oestrogen which can cause concern to livestock breeders. Freshly grazed forage causes most concern but the problem can be avoided by moving breeding animals off red clover around conception. Cattle are not normally affected but ewes should be taken off red clover at least a month either side of tupping.

#### Varieties

Modern plant breeding programmes have increased disease and pest resistance and improved persistence with varieties such as Milvus and Global.

There are two distinct types of red clover: early and late flowering. The former starts spring growth earlier in May followed by another growth flush. The latter flowers 10-14 days later after its one main growth period.

### 3 Alsike Clover (*Trifolium hybridum*)

A perennial which is slower to grow in the spring than red clover and is slightly lower yielding but otherwise has similar characteristics. Good for heavy and acidic soils.

### 4 Crimson Clover (*Trifolium incarnatum*)

An annual which can be sown after an early harvested cereal to provide winter sheep keep. It can also be used to give soil a fertility boost in a short period of time.

### 5 Persian Clover (*Trifolium resupinatum*)

An annual used to provide a quick boost to soil fertility on most soil types. It provides a good forage which may be grazed or conserved.

### 6 Berseem Clover (*Trifolium alexandrinum*)

Also known as Egyptian Clover, this is a short term, fast growing annual clover, which quickly provides large amounts of biomass and improves soil fertility. The least winter hardy of the true clovers.

## Other Key Legumes

### 7 Lucerne (*Medicago sativa*)

No one can really understand why so little lucerne (or alfalfa) is grown in the UK, when worldwide there are 13 million hectares cropped for forage. There are however a small number of UK farms now retrying this capable legume. Cut three times a year, it produces a protein-rich 14t DM per hectare without nitrogen fertiliser and on dry land.

Lucerne is a large plant with a similar erect growth habit to red clover. It is deep rooting, very drought resistant and has a yield high enough to be grown on its own. However it is usually sown with a companion grass such as meadow fescue or timothy which fill in the bottom of the crop.

Lucerne is useful to dairy farmers wanting to produce a high protein silage that is complementary to maize. It can be quite slow to establish and is only suitable for free-draining land that is not acidic.

### 8 Sainfoin (*Onobrychis viciifolia*)

Along with other forage legumes, sainfoin offers free nitrogen and extra protein content. But it has other benefits that mark it out as unique.

Sainfoin is capable of growing on the thinnest of alkaline soils, particularly the dry chalk and limestone land in the south of England. It is extremely drought-resistant and never stops growing, even in prolonged dry spells. Its root structure leaves



soil in excellent condition and sainfoin can be considered an invaluable part of a light land rotation. It penetrates soil and rock to a great depth where it seems able to extract nutrients better than any other species.

#### Boosting livestock production and health

Sainfoin contains tannins which aid protein absorption resulting in faster liveweight gains when compared to any other forage. This may also help reduce the amount of methane produced by ruminants, very useful from an environmental perspective. These tannins have another benefit: they mean sainfoin never causes bloat. Trials have shown that as little as 20% of sainfoin in the diet can offset the risk of bloat to near zero.

Sainfoin has a remarkable effect on wormy lambs, being a natural anthelmintic. EU projects 'Healthy Hay' and 'LegumePlus' have confirmed that feeding sainfoin disrupts the lifecycle of parasitic worms, so improving livestock performance yet further.

#### 9 Sweet Clover (*Melilotus* spp.)

Also known as yellow blossom, this biennial which has a feed value similar to lucerne can produce huge quantities of green material in July if sown in May. It is also a very good green manure, fixing a great deal of nitrogen and adding huge amounts of organic matter to the soil.

#### 10 Yellow Trefoil (*Medicago lupulina*)

This is a low growing, short-lived plant which sheds seeds freely and so regenerates itself. It is sometimes included in seed mixtures to give early spring growth which is unusual as most legumes are quite late to start growing.

#### 11 Birdsfoot Trefoil (*Lotus corniculatus*)

Like sainfoin, this legume contains tannins and is best suited to poorer soils where it outperforms other legumes. Including birdsfoot trefoil in seed mixes may offer other medicinal benefits, something that is currently being researched.

#### 12 Vetches (*Vicia sativa*)

This legume, also known as tares, when sown in the autumn or spring can provide one large crop for silage, and is excellent at out-competing weeds, fixing large amounts of nitrogen and improving soil structure.

## Herbs

Deep-rooting herbal leys are becoming popular on many farms as they offer huge benefits to livestock and soil structure. Using deep-penetrating roots instead of diesel-consuming tractors, herbal leys are an alternative way to aerate soil.

Agricultural herbs also provide minerals, essential for normal, healthy animal growth. Single species grass swards are often found to be lacking in these micro-nutrients. Deep-rooting herbs are a rich source of these and are currently being researched by agricultural scientists. Many expert farmers consider that adding these valuable plants to seed mixtures is a logical step.

#### 13 Chicory (*Chicorium intybus*)

A true 'ground breaking' plant with deep roots that can penetrate plough pans and grow well on the driest soil. This high-yielding perennial is a rich source of minerals and has althelmintic effects. It is therefore excellent for sheep or cattle threatened by intestinal parasites.

#### 14 Ribgrass (*Plantago lanceolata*)

This reliable perennial herb, also known as ribwort plantain, is relatively low yielding but has deep roots and is grown for its vitamin and mineral content (especially copper, calcium and selenium).

#### 15 Yarrow (*Achillea millefolium*)


Yarrow is a deep-rooting perennial and a rich source of vitamin A.

#### 16 Burnet (*Sanguisorba minor*)

On light, alkaline soils this is a long lived perennial forage. All parts of the plant are palatable and it is extremely drought resistant.

#### 17 Sheeps Parsley (*Petroselinium crispum*)

A short lived but useful herb, suits lighter soil types.







# Over-Seeding

Over-seeding is a simple, effective and low cost way to improve worn leys or old pasture without ploughing or reseeding.

## Sowing and Growing

### Suitable soils and optimum pH

Over-seeding can be beneficial on most soil types. Routine monitoring of pH levels will allow for any necessary corrections to be made.

### When to sow

When soil temperatures are above 7°C, usually between March and September. Sufficient soil moisture is vital. Avoid seeding into competitive swards during May and June when excessive grass growth will smother new seedlings.

### How to sow

Broadcast or shallow drill into recently grazed or cut leys. Before sowing, create a tilth using a chain or comb harrow. After sowing, roll thoroughly using a ring or flat roller, or tread in with sheep. Grass drills such as the 'Moore's' or 'Aitchinson' can be used. Cereal drills should be avoided as they can sow the grass seed too deep. Clover should never be sown deeper than 1cm. Spinners such as the 'Stocks' are good for applying small quantities of clover.

### Management

Gentle grazing should be resumed around five weeks after sowing. Cattle or sheep may be employed, but sheep should not be left on for long as they will graze too close, damaging new seedlings. Although cattle exert more pressure on the ground, they do not bite so accurately or as close and are the preferred choice provided that dry ground conditions prevail.

### Nutrient requirements

N fertiliser applications should be delayed until the new grass or clover seedlings are well established and able to tolerate the competitive growth that fertiliser brings. P and K levels should be maintained at ADAS Index 2.

Around a quarter of the grass seed sold in the UK is used for over-seeding. This seed is sown to improve worn or damaged leys and for patching up recent sowings which have not taken well.

To many farmers, over-seeding has advantages over the plough. It's cheap, quick and low risk, with existing grass being retained and improved without loss of forage or time.

Not all grasses and clovers are suitable for over-seeding. The best results come from the large seeded and vigorous strains of tetraploid ryegrass.

**The best results come from the large seeded and vigorous strains of tetraploid ryegrass.**

Of these, the Italian and hybrid forms are the quickest and best for cutting, with perennials being ideal for grazing leys.

White clovers usually give good results when sown into warm, moist soils especially where careful post-sowing grazing management is practiced.





Grass

Legume

Herb

## Mixes: Ryegrass

## Ryegrass Over-Seeding

Short Term 2-3 Years

Code: MIXOS

Ideal for the short term improvement of silage leys. The mixture is very competitive and provides good early spring growth. First cut is usually taken between the second and third weeks of May.

- 7.00 kg certified BARMULTRA II tet. Italian ryegrass
- 3.00 kg certified KIRIAL hybrid ryegrass

**10.00 kg/acre - £32.55**      25.00 kg/ha - £81.38

## Ryegrass Over-Seeding

Longer Term 4-5 Years

Code: MIXOSL

A flexible mixture for grazing or cutting fields which require longer term improvement. The grasses will provide growth from spring through the summer.

- 5.00 kg certified NOVIAL hybrid ryegrass
- 5.00 kg certified CALIBRA tet. perennial ryegrass

**10.00 kg/acre - £40.75**      25.00 kg/ha - £101.88

## Ryegrass &amp; Clover Over-Seeding

Longer Term 4-5 Years

Code: MIXOSLC

A combination of ryegrasses and a half-rate of persistent clovers, this mixture can be grazed by sheep or cattle and can also be cut for silage.

- 4.50 kg certified NOVIAL hybrid ryegrass
- 4.50 kg certified CALIBRA tet. perennial ryegrass
- 0.80 kg certified ABERDAI white clover
- 0.20 kg certified ABERACE wild white clover

**10.00 kg/acre - £47.54**      25.00 kg/ha - £118.85

## Additions



## Bottom grass

The addition of a bottom or grazing-type ryegrass can help to fill in the sward in open leys.

Add 2kg of grazing ryegrass

**£10.50 per acre**

## Mixes: Clover and herbs

## White Clover Over-Seeding

Long Term Grazing

Code: MIXOSC

This persistent mixture combines medium and small leaved clovers which provide grazing for sheep or cattle. It may also be used for silage making.

- 0.80 kg certified ABERSWAN white clover
- 0.80 kg certified ABERDAI white clover
- 0.40 kg certified ABERACE wild white clover

**2.00 kg/acre - £21.72**      5.00 kg/ha - £54.30

## White Clover Over-Seeding

Dairy Graze or Silage

Code: MIXOSCD

Using highly productive medium and large leaved white clovers this mixture is ideal for dairy grazing or silage making. It can also be grazed by sheep occasionally if required.

- 1.00 kg certified ABERDAI white clover
- 1.00 kg certified ALICE white clover

**2.00 kg/acre - £21.30**      5.00 kg/ha - £53.25

## Herbal Over-Seeding

Deep-Rooting Herbal ley

Code: MIXHOS

Deep rooting herbal leys are becoming more and more popular. Grass-only swards lack protein rich clovers and mineral rich herbs. Ideally, herb-rich swards are best established by reseeding but where this is not possible this mixture can be oversown into a grass-only sward.

- 0.40 kg certified LEO birdsfoot trefoil
- 0.35 kg certified GLOBAL red clover
- 0.25 kg certified ERMO alsike clover
- 0.25 kg certified MERWI white clover
- 0.20 kg certified ABERDAI white clover
- 0.10 kg certified ABERACE wild white clover
- 0.20 kg commercial sweet clover
- 2.00 kg commercial sainfoin
- 1.00 kg burnet
- 0.50 kg sheeps parsley
- 0.35 kg certified LACERTA chicory
- 0.25 kg certified ENDURANCE ribgrass
- 0.15 kg yarrow

**6.00 kg/acre - £54.28**      15.00 kg/ha - £135.70



# Hay Leys

Grass only hay leys that offer high quality and bulk.

Grass



## Sowing and Growing

### Suitable soils and optimum pH

The ryegrass based leys are best suited to fertile and moisture retentive soils. Ryegrass can suffer on drought prone soils, so an option on dry land could be to add a deeper rooting, inexpensive festulolium plant to the mixture - please enquire when ordering.

### When to sow

The crop should be sown August-September to provide good yields the following spring. Except for the westerwold hay mix, these mixtures won't put on a seedhead when sown in the spring.

### How to sow

A non-selective herbicide should be used before seedbed preparation. Drill into a fine firm seedbed and try to avoid drying the soils out with excessive cultivations in dry autumns. Rolling to retain moisture and break down clods before and after sowing with a cambridge or flat roller is essential. Broadcast seed should be harrowed lightly after sowing and before rolling.

### Management

Annual weeds should disappear as the new seeds begin to take over, or they can be grazed out with stock. Cutting usually begins in late June and takes place before and during flowering. Graze excess growth after the required cut has been taken by November to avoid winter kill.

### Nutrient requirements

These leys should receive approximately 70kg ha of nitrogen. Excessive applications of nitrogen can cause the plant to become sappy and difficult to dry. As cutting and removing depletes P and K levels, they should be maintained around ADAS index 2.

## Mixes

### Hard Horse Hay

#### Two Year Ley

Code: MIX9

Devised specifically for the production of hay or haylage. With good disease resistance it produces a consistent sample of hard hay. Although grazable, it's principally a cutting ley. Sow in autumn to provide stemmy hay the following spring.

- 8.00 kg certified BARMULTRA II tet. Italian ryegrass
- 3.00 kg certified JAVORIO Italian ryegrass
- 3.00 kg certified SHAKIRA Italian ryegrass

**14.00 kg/acre - £42.95**

35.00 kg/ha - £107.38

### Hay and Graze

#### Four Year Hay/Haylage Ley

Code: MIXHG

A longer term option for the hay & haylage producer with upright hybrid ryegrass and longer lasting perennial ryegrass. Diploid varieties are included for faster drying. Sow in autumn to provide a crop the following spring, or cut earlier in the year for haylage. Also provides high quality summer & autumn grazing.

- 5.00 kg certified BARCLAMP dip. hybrid ryegrass
- 6.00 kg certified ABERWOLF perennial ryegrass
- 2.00 kg certified COMER timothy

**13.00 kg/acre - £59.95**

32.50 kg/ha - £149.88

### Traditional Hay Maker

#### Long Term Hay Ley

Code: MIXHM

Slower to establish than a straight ryegrass ley but will provide good quality, bulky hay crops with low disease levels for many years. Once cut it can be used for aftermath grazing.

- 5.00 kg certified ABERWOLF perennial ryegrass
- 5.00 kg certified PARDUS meadow fescue
- 3.00 kg certified WINNETOU timothy

**13.00 kg/acre - £71.80**

32.50 kg/ha - £179.50

## Mixes

### Westerwold Hay Mix

#### One Year Ley

Code: MIXWWH

This flexible cutting option, produces a short term, clean, high quality hay, due to its rapid growth and high biomass it can be difficult to dry properly. This mix has been designed to include high levels of diploid westerwold which contains less moisture and dries evenly, speeding up the hay making process.

- 10.50 kg certified LIFLORIA dip. westerwold ryegrass
- 3.50 kg certified JIVET westerwold ryegrass

**14.00 kg/acre - £39.90**

35.00 kg/ha - £99.75

## Additions



### Sweet vernal grass

To create a softer, sweeter smelling meadow hay.

Add 0.1 kg of sweet vernal grass

**£6.00 per acre**





# Silage

Good silage comes from a good ley.

Good silage depends on many factors. These include appropriate fertiliser applications, growth stage when cut and how the crop is wilted and stored. But the most important factor is to select the right crop species and varieties to suit the soil type from the start.

## Ryegrass leys

Ryegrass in all its forms (see page 4) has been the building block of short term silage leys for the last 60 years. With the various high yielding types such as westerwolds, Italian, hybrid and perennial lasting between one and five years, there is a ryegrass variety to suit every system. Highly responsive to nitrogen, ryegrass-based swards produce palatable silage that increases milk and meat production.

## Red clover leys

With its high yields, forage quality and suitability for silage, red clover swards are playing an increasingly important role in sustainable systems of grassland farming, especially now nitrogen prices are so high.

At 19% crude protein, red clover's nutritional value is higher than grass' and its high voluntary intake leads to enhanced animal performance.

Thriving on most soils, its ability to 'fix' atmospheric nitrogen in the root nodules (an average of 200kg N/ha), offers a saving on manufactured nitrogen fertiliser.

Red clover is tolerant to winter cold and, due to its deep rooting characteristic, is drought resistant. Used as a break crop it will improve soil structure and fertility while also giving excellent forage yields.

## Lucerne

At 20% protein lucerne is an attractive feed. It is a good complement to maize and is leafy and low in fibre, breaking down rapidly in the rumen and passing out quickly, allowing a greater intake of forage than many other species. Lucerne has significant benefits but few people grow it believing, incorrectly, that it is a difficult crop to maintain.

## Sainfoin

Sainfoin performs better than any other crop on thin, dry, calcareous and brashy soils. This remarkable plant is extremely drought resistant with its deep-penetrating roots. In addition, it needs no nitrogen fertiliser and very little phosphate. It offers a protein-rich forage with medicinal qualities that will appeal to all types of livestock farmer.

## Vetch

This is a short term annual with a high protein and mineral content. Vetch is fast to grow and can be sown alone for silage or grazing and is also suitable for mixing with cereals such as oats for whole-crop silage. Quick to establish, it can also be sown with grass and clover mixes to produce extra yield.

## Great Silage, Great Soil

Short term leys are beneficial in arable rotations and are a solution on the many farms with deteriorating soil structure.

Ryegrass leys produce a large amount of root mass in a short time which improves soil structure when it decays at the end of the ley's term. Deep-rooting legume-based leys are also excellent at improving soil, and have the additional benefit of fixing nitrogen, invaluable at a time of rising fertiliser prices.

These leys are also effective in the battle against blackgrass as a one, two or three year ley breaks the lifecycle of this weed, so benefiting subsequent crops.





# Intensive Silage

Short and medium term leys that provide the highest yields for silage.

In these leys the various types of ryegrass (see page 4) have been combined to produce high-yielding quality silage crops lasting between one and five years. The characteristics of individual varieties have an impact on the timing of the first cut and the potential number of cuts per year. There is also the potential to graze the aftermath with many of these mixtures.

## One year bulk

There has been a sharp increase in the use of Westerwolds. Rapid in establishment and quick to produce bulk, these leys are very useful for producing early grazing and silage from autumn sowings. They can also be sown in the spring on moisture-retentive soils to provide heavy summer silage crops.

Unlike other ryegrasses, westerwolds will produce a stem and seed head from a spring sowing. This is useful, especially for the production of high quality horse hay.

## High yield, high quality

Our two to five year seed mixtures are formulated for the focused and intensive farmer who requires silage and grazing leys to have a direct impact on milk or meat production.

In these economic times, the need to maximise milk and meat from efficient forage production is clear. Our mixes are therefore designed to combine exceptionally good yields with the highest nutritive value.

Emphasis is placed on achieving optimum D-value at the time of utilisation, as well as high soluble carbohydrate content. Grass varieties in these mixtures give ultimate performance and are highly rated for overall disease resistance which will improve both yield of grass and palatability of forage.

## What is festulolium?

Festulolium is a natural hybridisation of ryegrass and fescue species, combining the stress resistant genes of fescue with the bulky yield of ryegrass. It provides a more resilient species, with a better tolerance of drought or water logged soils, while still providing high yielding, very palatable forage. Festulolium is generally an upright grass, suitable for cutting.

## Sowing and Growing

### Suitable soils and optimum pH

Ryegrass is best suited to fertile and moisture-retentive soils and will tolerate slightly acidic pHs (6.2-6.5).

### When to sow

Highest yields come from autumn sowings (August – September). Spring sown crops (March – May) are dependent on sufficient moisture and should be avoided in drought-prone areas.

### How to sow

Drill in two directions into a fine, firm seedbed at 10-20mm. Rolling with either a Cambridge or flat roller before and after sowing is recommended. Broadcast seed should be harrowed lightly after sowing and before rolling.

### Management

When spring sown these leys can be ready for cutting after only 12 weeks. A further two cuts can follow on soils that have adequate moisture and N. Early cuts just prior to heading will give high D-values and good regrowth. Cut frequently to encourage high D-value leafy growth. Graze excess growth by November to avoid winter kill.

### Nutrient requirements

100kg N/ha for the first cut with 60-80 kg of N applied per subsequent cut. As cutting depletes P and K levels, these should be maintained at ADAS Index 2.

### Yield potential

Westerwolds:	18t DM/ha
Italian ryegrass:	18t DM/ha
Hybrid ryegrass:	14t DM/ha
Perennial ryegrass:	13t DM/ha

Typical silage analysis has a dry matter of 30%, a crude protein of 16%, a D-value of 70 and an ME of 11MJ.



Grass

Legume

## Mixes

## Quick Bulk Westerwolds

## Intensive One Year Ley

Code: MIXQB

A very fast growing ley which can be sown in the spring or autumn and is primarily grown to produce silage. However, it can be grazed and the crop can provide 'early bite' when sown in the autumn. It is therefore a cheap alternative to cereal rye which was formally grown for this purpose.

- 10.00 kg certified JIVET westerwolds ryegrass
- 4.00 kg certified LIFLORIA dip. westerwolds ryegrass

**14.00 kg/acre - £39.90**    35.00 kg/ha - £99.75

## Festulolium Silage Ley

## Two Year Dry Land Ley

Code: MIXAF

This mix contains one of the festulolium varieties on the UK recommended list. It is a hybridisation of Italian ryegrass and meadow fescue, combining the bulky yield of Italian ryegrass with the stress resistant genes of fescue. An ideal mixture for dryland, it showed good resilience during the dry summer 2018. Both species will head in the third week of May and can be relied on for 2 years.

- 10.00 kg certified ABERNICHE festulolium
- 4.00 kg certified BARMULTRA II tet. Italian ryegrass

**14.00 kg/acre - £53.50**    35.00 kg/ha - £133.75

## Westerwold and Vetch

## Six Month Ley

Code: MIXWWV

A good balance between a vigorous grass and a fast growing short term legume, this mixture can be used to provide a very large cut or early spring grazing. As westerwolds will regrow after cutting this ley can be left for a further cut or grazed if required. To minimise the risk of ryegrass seed being shed, it is advisable to cut before the seed heads are visible.

- 8.00 kg certified JIVET westerwolds ryegrass
- 17.00 kg certified EARLY ENGLISH vetch

**25.00 kg/acre - £53.40**    62.50 kg/ha - £133.50

## Additions



## White Clover

White clover can be added to improve nutritional value.  
Add 1kg of white clover    **£10.90 per acre**

## Festulolium

Swap ryegrass for festulolium for improved drought tolerance.  
**Please call for advice**

## Mixes

## Maximum-Yield

## Two Year Silage Ley

Code: MIXA

Optimum balance between the highest quality and yield for silage, Maximum-Yield produces the all-important first cut between the second and third week of May. At this time the grasses will have high D-values and soluble carbohydrate content which ensures good silage fermentation.

- 10.50 kg certified BARMULTRA II tet. Italian ryegrass
- 3.50 kg certified JAVORIO Italian ryegrass

**14.00 kg/acre - £42.95**    35.00 kg/ha - £107.38

## Hybrid Silage Ley

## Three - Four Year Ley

Code: MIXB

Early growth, high yields and good persistence make hybrid ryegrasses worth considering when the ley is expected to remain down for more than two years. Utilising excellent hybrid varieties, this ley is comparable, in terms of output, with Italian ryegrass. It should also be noted that the grazing potential of this ley is superior.

- 6.00 kg certified KIRIAL hybrid ryegrass
- 6.00 kg certified NOVIAL hybrid ryegrass
- 2.00 kg certified ASTONKING perennial ryegrass

**14.00 kg/acre - £57.90**    35.00 kg/ha - £144.75

## Maximum D-Value

## Four - Five Year Silage Ley

Code: MIXC

Maximum feed value can only be obtained from well made high D value silage. This ley will provide optimum digestibility and yield during the third week of May. Yields are boosted by utilising hybrid ryegrass with high quality Aberwolf perennial ryegrass, noted for its very good D-value on the recommended list and providing good summer and late season grazing.

- 4.00 kg certified ABERWOLF perennial ryegrass
- 6.00 kg certified ABERCLYDE tet. perennial ryegrass
- 2.00 kg certified KIRIAL hybrid ryegrass
- 2.00 kg certified NOVIAL hybrid ryegrass

**14.00 kg/acre - £66.60**    35.00 kg/ha - £166.50





# Red Clover Leys

Red clover leys produce a protein rich 15t DM per hectare without nitrogen fertiliser.

Red clover produces silage with a 2-3% higher protein content than a grass-only equivalent. This, combined with its high intake characteristics, leads to improved milk and meat production.

Red clover is drought tolerant and like many fast-growing legumes it's able to 'fix' up to 250 kg N/ha. To provide enough free nitrogen for a successful crop, legumes need to be included at high proportions in a mixed sward.

Legumes do not fix nitrogen all year round. For this natural chemistry to occur, the soil needs to be warm and, in the UK, this usually means that nitrogen fixation occurs between April and September.

## With or without grass?

Red clover can be sown as a monoculture at 5-6 kg/acre for silage, but a mixture with grasses is preferable since this gives higher total forage yield and makes better silage. Mixtures of 9 kg/acre grasses and 3 kg/acre red clover are commonly sown to provide the correct balance. A pure stand of red clover generally yields lower than the grass and clover mixture at about 5-6t DM/ha.

For a one or two year ley Italian ryegrass is an excellent component, but for a duration of three years or more a mixture of hybrid and perennial ryegrass is a better option.

To allow full expression of the red clover, it is best to use tetraploid varieties of ryegrass since they tiller less densely than diploids. Their early-season ear emergence patterns should also coincide with the flowering pattern of the red clover. They are then at the same maturity stage and digestibility is similar.

## What you need to know about oestrogen

There are questions over the effect that the oestrogen content of red clover may have on reducing animal fertility. There are relatively few confirmed cases and it is commonly accepted because a ewes diet may be made up solely of red clover, it is best to flush and tup ewes on leys that do not contain red clover, do not feed or graze ewes 6 weeks before or after tugging to be safe.

There is no known detrimental effects on fattening lambs, in fact they can fatten very well on this high protein crop.

## Sowing and Growing

### Suitable soils and optimum pH

Grows on most soils, including the drought prone. The optimum pH is 6.0-6.5 for N fixation, but red clover will tolerate 5.6.

### When to sow

Sow from March until September. Red clover mixtures can be undersown in an arable crop, or after harvest provided there is enough time for the plants to develop sufficiently prior to winter cold.

On light soils in dry districts autumn sowings perform better as these will have well established roots capable of better growth in dry seasons.

### How to sow

For sound establishment, a well cultivated, firm, level seedbed is needed to ensure that the small clover seeds are drilled uniformly at a shallow depth of 10-15 mm. The use of a roller prior to and after sowing is essential.

### Management

The competitiveness of red clover against weeds is low at the early establishment phase particularly if sown alone.

Topping is of value and 'clover-safe' herbicides are available, though they can check red clover development to some degree. To avoid clover sickness (a combination of soil-borne sclerotinia and stem eelworm) a five year gap should be allowed between leys containing red clover.

### Nutrient requirements

Red clover will fix its own N, but P and K levels must be maintained at an ADAS Index 2. A small quantity of N can be applied in the autumn or early spring to enhance initial growth. This should not exceed 25kg N/ha.

### Yield potential

Forage yield in the establishment year of a spring-sown sward is circa 60% of that possible in the first harvest year which should be around 15t DM/ha.

The yield is spread over 2-3 cuts per year. Typical silage analysis has a dry matter of 30%, a crude protein of 19%, a D-value of 72 and an ME of 12MJ.



Grass

Legume

## Mixes

## Fast and Vast

One - Two Year Ley

Code: MIXFV

This short term ley is for those wishing to produce a large amount of forage with little or no nitrogen fertiliser. Yields are high, especially on rich, moist soils and the majority of crops are made into silage. In addition to red clover, the mixture also contains crimson clover and vetch which increase yield over a short period of time. It can be relied upon for one full year of production or left down for a second.

- 10.00 kg certified EARLY ENGLISH vetch
- 2.00 kg certified AVISTO red clover
- 1.00 kg certified HEUSERS OSTSAAT crimson clover
- 4.50 kg certified FABIO tet. Italian ryegrass
- 4.50 kg certified SHAKIRA Italian ryegrass

**22.00 kg/acre - £67.65**

55.00 kg/ha - £169.13

## Short Term Red Clover Ley

One - Two Year Mixture

Code: MIXCG03

An intensive silage ley lasting for up to two years which requires little or no nitrogen fertiliser. First cut is to be expected during the third week of May.

- 3.00 kg certified AVISTO red clover
- 3.00 kg certified BARMULTRA II tet. Italian ryegrass
- 3.00 kg certified FABIO tet. Italian ryegrass
- 3.00 kg certified JAVORIO Italian ryegrass

**12.00 kg/acre - £52.80**

30.00 kg/ha - £132.00

## Longer Term Red Clover Ley

Three - Four Year Mixture

Code: MIXCG06

Persistent and high yielding, this ley is tried, tested and highly successful. It is usually cut in the third or fourth week of May and incorporates the best red clover with hybrid and perennial ryegrasses, giving yields nearly as high as our two year red clover ley.

- 3.00 kg certified MILVUS red clover
- 3.00 kg certified KIRIAL hybrid ryegrass
- 3.00 kg certified NOVIAL hybrid ryegrass
- 3.00 kg certified CALIBRA tet. perennial ryegrass

**12.00 kg/acre - £62.70**

30.00 kg/ha - £156.75



**Red Clover**  
Gloucestershire  
28th May

## Additions

**Vetch**

Vetch may be added to red clover and ryegrass mixes to increase yield in the first growing season.

Add 10kg of vetch

**£18.00 per acre**



# Sainfoin

High yielding silage or hay crop with occasional grazing for dry, alkaline soils. Bloat free and a natural anthelmintic.

## Sowing and Growing

### Suitable soils and optimum pH

Performs best on free-draining alkaline soils. Do not sow on land below 6.2pH.

### When to sow

Always sow sainfoin into warm soils in the spring.

### How to sow

Sainfoin seed can be undersown to spring cereals or direct drilled in April or May at around 30mm. If undersown, the cereal sowing rate should be reduced to 40 kg/acre.

### Management

A sainfoin ley should be managed carefully to maximise performance. Sainfoin produces a cut of silage in early June or hay may be taken if preferred. Sainfoin should be cut during early flowering but this may be delayed without much loss of feed value if needed. Regrowth is less after the first cut and may be cut again or grazed. Grazing should be light and quick to avoid damage to the plant. Never set stock it or it will become thin.

### Nutrient requirements

Sainfoin requires no N or P but K levels must be maintained at ADAS Index 2 to safeguard yields.

### Yield potential

14t DM/ha annually. Typical silage analysis has a dry matter of 14%, a crude protein of 18%, a D-value of 62 and an ME of 9.5 MJ. However, sainfoin produces better results than this analysis indicates as its high tannin content protects the protein in the rumen so increasing absorption and producing higher liveweight gains.



There are few crops quite like sainfoin. It is a high-yielding, drought-resistant plant which needs no nitrogen fertiliser and little phosphate. It won't cause bloat, is a natural anthelmintic and, with rumen-protected protein, produces top quality meat and milk.

Sainfoin has deep-penetrating roots making it highly suitable for the dry, alkaline soils of England. In these times of fluctuating fertiliser, feed and veterinary drug prices, alongside increased demands for sustainability, there are few crops that tick as many boxes.

It grows best on stony brash or chalks, but does not like wet soils where red clover should be chosen in preference.

## Sainfoin

### Four Year Cutting or Grazing

Code: SAI

On the right ground this is a superb crop. Lasting for four years or more, it is extremely valuable for finishing lambs.

■ 35.00 kg commercial sainfoin

**35.00 kg/acre - £98.00**

87.50 kg/ha - £245.00

## Companion Grass Option

### Four Year Mixture


Code: MIXLUC

We recommend the use of a non-competitive grass mixture to be sown with sainfoin. The grass fills the base of the crop, increasing yield and soluble sugars to improve silage fermentation. The grass seed element should be surface sown and rolled in.

■ 2.00 kg certified PARDUS meadow fescue  
■ 1.00 kg certified WINNETOU timothy

**3.00 kg/acre - £17.15**

7.50 kg/ha - £42.88



For more information on sainfoin, please download our growers guide - **Sainfoin - Surprising science behind a forgotten forage.**

Visit [cotswoldseeds.com](http://cotswoldseeds.com) to download your copy.



**Sainfoin**  
Wiltshire  
29th May



Grass

Legume

# Lucerne

Reliable yields for silage on dry gravels.

Lucerne is highly productive and reliably provides three to four cuts of protein-rich silage annually, even through drought, and lasts for around five years. Lucerne must be grown on naturally alkaline and free draining soils or gravel. It is slower to establish than ryegrass and does require careful management but, if agronomic guidelines are followed, there is nothing complicated about it.



## Lucerne

### Four Year Cutting Crop

Code: MARSH

Lucerne should be sown as a four or five year temporary ley. For specialist use it may be made into hay for the equine market where it is known as alfalfa.

■ 8.00 kg certified TIMBALE lucerne

**8.00 kg/acre - £66.00**

20.00 kg/ha - £165.00

## Companion Grass Option

### Four Year Mixture

Code: MIXLUC

We recommend the use of a non-competitive grass mixture to be sown with lucerne. The grass fills the base of the crop, increasing yield and soluble sugars to improve silage fermentation. The grass seed element should be surface sown and rolled in.

■ 2.00 kg certified PARDUS meadow fescue  
■ 1.00 kg certified WINNETOU timothy

**3.00 kg/acre - £17.15**

7.50 kg/ha - £42.88

## Sowing and Growing

### Suitable soils and optimum pH

Gravels and free-draining soils with a pH 6.5-8.

### When to sow

Lucerne must be sown into warm soils and is often undersown to a spring cereal crop as it is slow to establish. Reducing the cereal seed rate by a third and cutting it as arable or wholecrop silage will give lucerne the best start. Alternatively, sow in the summer following an early-harvested cereal such as winter barley. The middle of August is the latest date for sowing.

### How to sow

The seed of lucerne is small and needs to be sown to a maximum depth of 15mm otherwise a patchy, thin crop will result. Roll before and after sowing to help achieve fast germination and weed competitiveness. Sowing with a companion grass mix helps out-compete weeds enabling lucerne leys to be left down for longer. Herbicides exist but are limited and you should obtain a clean seedbed prior to sowing.

### Management

Following a direct spring sowing a light cut may be taken in mid August. From a summer sowing or an undersowing there will be little to cut in the first year. Leave until the following spring when it should be cut for the first time in early June at almost full flowering. Thereafter cut at the bud stage as this provides the ideal balance between yield and quality. Two or three further cuts follow at six week intervals. After cutting, the crop needs wilting so that it contains less than 70% moisture when made into baled silage. Hard or frequent grazing should be avoided especially during its first year as the crop will not tolerate it. Lucerne can also cause bloat when grazed.

### Nutrient requirements

Although lucerne requires no N once established it can be beneficial to apply 25kg N/ha to an autumn sowing to promote rapid plant development. P and K requirements are higher than for grass and should be maintained at ADAS Index 2 to maintain yields.

### Yield and nutrient data

14t DM/ha annually. A well fermented lucerne/grass silage has a dry matter of 30%, a crude protein of 20%, a D-value of 60 and an ME of 9.7MJ.

## Culture




### To Fix N

The use of culture to provide the correct type of bacteria to initiate nodulation is considered essential. Mix with seed on the day of sowing.

Sachet for 25 kg of seed

£10.00 plus VAT





# Grazing

Seed mixtures to suit the UK's many grazing systems.

Whether you want grass to grow in the uplands or the lowlands, on dry or wet soil, on acidic, neutral or alkaline soils, we can provide a seed mix to suit.

The number of species in our grazing mixtures varies from one to eighteen, providing a huge range of choices to meet the requirements of the diverse grazing systems on farms across the country.

Pioneers of grassland management such as Andre Pochon, Robert H Elliot and William Lamin developed complex mixtures of grasses and legumes for grazing and cutting. In recent times however, intensively bred strains of ryegrass in temporary leys have been very popular as they respond well to nitrogen fertiliser and give the high yields required by intensive farmers



## Clover versus N fertiliser

Nitrogen fertiliser is one of the largest costs to livestock farmers at around £200 per hectare. Now, in light of unpredictable fertiliser prices, many farmers are sowing high-clover leys to lower the cost of forage production as white clover and other nitrogen fixing legumes can reduce or replace the need for artificial nitrogen.

## Single species versus diversity

While some have good reason to grow single species swards, sowing mixtures of grasses and clovers offers real benefits. A single grass alone will often be lower yielding and more vulnerable to failure or poor performance due to pests, disease or the effects of unusual weather. A diverse mixture is therefore more reliable and preferable to sowing a single species.

This is especially important for leys which are expected to last for more than one year. Higher yields from mixtures of grasses and clovers are due to better seasonal distribution of growth: grasses give high yields during May and June, clovers produce theirs in July and August. Critically, it's the contribution of both grass and clover that provides the optimum balance between bulk yield and feed value. Grasses tend to have higher annual yields, but are lower in protein than clovers. Animals grow faster and do better on a mix of clover & grass.

## Alternatives to ryegrass

Timothy and meadow fescue are generally considered to be the most palatable of the permanent grasses. Although they may lack some of the digestibility of ryegrass, they are consumed readily by the grazing animal. In addition, when grown with red and white clovers, the forage produced will be higher in protein, more digestible and largely self-sufficient with little demand for artificial nitrogen. They also offer impressive yields. These grasses are excellent in mixtures and a very good alternative in circumstances where ryegrass doesn't suit, such as low fertility and/or wet soils or in the uplands.

## Drought resistant swards

In recent summers extended dry periods have put a real strain on livestock farmers battling to ensure they have sufficient forage year round. Grass species such as cocksfoot and clover continue to produce even when there has been no rain for weeks, and many of our mixes are designed with these conditions in mind.

## Herbal Leys: feeding health

The most diverse grazing mix we offer is the herbal ley which contains a huge range of grasses, herbs and clovers. It produces well-balanced forage, not just large volumes of grass, and thrives in dry conditions. Species such as cocksfoot, red clover and chicory are deep-rooting soil improvers with the ability to unlock mineral resources from deep in the soil profile.

Herbs are richer in minerals than grasses or clovers and including them in seed mixes is an effective way of improving forage to ensure good animal health and performance.

## Yield and longevity

A newly sown ley on good soil, with plenty of moisture and nitrogen will significantly out-yield older swards. Over time, deterioration of any seed mix is inevitable as unsown, less nutritious species invade. Mixes containing late heading ryegrasses (such as Cancan) have greater persistence, so reducing the need to reseed frequently. To maximise a ryegrass-based ley's productive lifespan, nitrogen levels should be maintained as a drop in fertility will reduce the competitiveness of ryegrass, so favouring unsown species.



First Hand

## Henry Edmunds



<b>Farm Type</b>	<b>Mixed</b>
<b>Location</b>	<b>Hampshire</b>
<b>Size</b>	<b>2500 acres</b>
<b>Soil Type</b>	<b>Light, calcareous</b>
<b>Mixes Used</b>	<b>Bespoke Cholderton ley</b>

Henry Edmunds advocates using what he describes as an 'integrated regenerative ley farming system' to get the most from poor soil quality, manage blackgrass, feed his stock and provide habitats for wildlife. At the heart of this ley farming system is the bespoke, herb rich, Cholderton ley, named after a mix devised by Henry.

The Cholderton Estate, Henry's family farm, lies on the fringes of the Salisbury Plain, where he runs a mixed operation with a spring-calving dairy herd, beef cattle, plus a flock of Hampshire Downs sheep. The soils are thin and overlie a bed of chalk and flint and the land is classed as Grade 4 with severe limitations for crops and yield.

'The poor quality of the land makes it more vulnerable to climate change, to heavy rains and prolonged periods of drought,' says Henry. 'But we mitigate this with the leguminous rich leys. The ley farming system builds fertility without the need for fertilisers or agro-chemicals. The sophisticated mixture of grass with legumes and herbs in the bespoke Cholderton Mixture produce higher yields and is also long-lasting.'

'I formulate the mixtures with deeper rooted plants - Timothy, meadow fescue, cocksfoot, red fescue and legumes including lucerne, sainfoin and clover mixtures,' Henry

explains. 'Lucerne is deep rooting and highly productive, out performing other forage species in yield, while sainfoin is drought-resistant, highly palatable, grows earlier than lucerne and is a natural anthelmintic. Stock grows faster on this than on any other forage crop. White clover fills the base of the sward with more high quality forage.'

'Meanwhile, below ground, the roots grow up to a metre down, capturing and locking in carbon. Nodules capture atmospheric nitrogen and the results are luscious growth of highly digestive forage. The mixture produces good yields on the poorest soils, while simultaneously enriching the soils.'

The leys can be sown in the autumn after the previous cereal crop has been harvested. Sheep should be encouraged to clear the weed burden in the late autumn or early winter. By the spring there will be a mixture of the cultivated species and blackgrass, but this will disappear over the following summer due to the grazing and cutting regime. The perennial grasses and legumes achieve a supremacy over the annual blackgrass, removing it by simple competition with the assistance of grazing.

### **"The mixture produces good yields on the poorest soils, while simultaneously enriching the soils"**

Blackgrass, being an annual, requires cultivation to germinate and under this management will disappear by the second or third year. Sheep and cattle will selectively graze out blackgrass; this together with a silage cut and the new system will eradicate this noxious weed.

After 5 years, the ley is ploughed in. This will produce a seed bed rich in organic matter and nutrients and will give an excellent crop of wheat with much lower inputs than would normally be required.

Every year, a quarter of the cereal crop is undersown. The grass and legume seeds are drilled into the newly sown cereal crop meaning the young grass and legumes are protected by the cereal, both from weeds and direct sunlight. In the autumn, after harvesting, the undersown grass emerges through the stubble and provides autumn grazing. The cereals are ground and fed to livestock over the winter with some retained as seed for the following year.

The herb rich leys are tailored for ELMS or CSS. 'It's wonderful to see arable fields reverting to downland and alive with rare plant species and insects,' says Henry.





# Dual Purpose Swards

Self-sufficient leys that provide high protein grazing with little or no nitrogen fertiliser.

## Sowing and Growing

### Suitable soils and optimum pH

These ryegrass-based leys grow on all but the most waterlogged soils. They are best suited to pH6 or above but will grow down to pH5.6.

### When to sow

Seed can be sown between March and mid September when soils are warm and sufficient moisture is available.

### How to sow

A very firm seed bed is required as white clover and timothy seeds are small and benefit from shallow sowing at around 10mm. Rolling at least once after sowing is strongly recommended. If under-sowing, do not choose a thick crop as this will result in failure; a thinly sown spring cereal is ideal. A clover seed mix is best sown by broadcasting with a spinner but can also be sown with a proper grass seed drill.

### Management

As the main period of grass growth is during May and June, it makes sense to take a cut of silage or hay during this time. Additionally, where grass growth exceeds grazing demand more can be cut and round-baled as silage.

Ideally, these leys should be rotationally grazed with an interval of 3-5 weeks for recovery and regrowth.

Seed mixtures containing hybrid ryegrass should be relied upon for up to four years only.

### Nutrient requirements

These leys should receive no more than 100kg N/ha with the biggest demand (60kg N/ha) in March and April if a silage cut is to be obtained. Pochon Dairy requires very little or no N as the high clover content (30-50%) will fix N in the soil.

### Yield potential

12t DM/ha should be achieved.

These mixtures are ideal for those looking to graze and/or cut a medium to long term ley. With white clover included in all three, they are largely self-sufficient in nitrogen. Of course the benefits derived from clover are proportional to the amount in the sward, both in terms of animal nutrition and nitrogen fixing.

If using the mix for silage or hay, the ley should be shut up at least six weeks before cutting, with the best combination of yield and quality coming from grasses that are just beginning to produce a seed head and clovers in bud or early flower. A first cut of silage is ready during late May. These leys will provide a second cut but are usually grazed.

These three mixes are among our most popular. The 'Milk-Meat' combination has been in use since the Second World War and sown on, we estimate, over a quarter of a million acres.

## The case for clover

Red clovers can be broken down into two distinct types: early and late flowering with a difference between flowering periods of 10 - 14 days.

Early or double cut red clover – These are most popular and are commonly used for silage as they re-grow well to provide a second cut later in the year.

The late varieties such as Atlaswede can be used on late, wet ground to provide one large single cut later in June.

### White clover

White clover is classified according to leaf size, which break down to small, medium and large, the size of the leaf dictates what type of use it is most suited to.

Small Leaf Clover - The very small, low yielding but extremely persistent type is known as 'wild white'. Little clovers like AberAce are extremely persistent, filling the base of the sward and can be grazed hard especially with sheep.

Medium Leaf - Medium leaved varieties such as AberSwan and AberDai offer yields well in excess of the wild whites. They are also more competitive, persistent and offer good early spring growth.

Large Leaf - The large leaf types such as Alice are the highest yielding. However, large leaved varieties do not survive well when grazed hard with sheep. Therefore these are best sown in silage or cattle grazing leys. For most situations it is best to sow a mixture of types to allow for grazing or silage.



Grass

Legume

## Mixes

## Early Bite

## Sheep and Hay Ley

Code: MIX1

Ideal for producing early grass on light land, this ley will provide good growth throughout the year and is especially good for lamb production. It is also suitable for hay or silage and can be expected to last for at least three years. Contains white clovers, making grazing more palatable and increasing live weight gains.

- 4.00 kg certified KIRIAL hybrid ryegrass
- 4.00 kg certified ABERWOLF perennial ryegrass
- 2.00 kg certified ASTONKING perennial ryegrass
- 1.40 kg certified WINNETOU timothy
- 0.20 kg certified ABERDAI white clover
- 0.20 kg certified ABERSWAN white clover
- 0.20 kg certified ABERACE wild white clover

12.00 kg/acre - £60.89

30.00 kg/ha - £152.23

## Pochon Dairy

## Two-Four Year Silage/Grazing Ley

Code: MIXCG02

Designed specifically for the dairy farmer wishing to produce silage and high quality grazing. This ley has an open growth habit allowing the white clover plenty of space to exploit. Including Aberystwyth ryegrass and white clovers, this mixture is principally intended to be grazed by the dairy cow. For sheep grazing use 'Pochon' Persistent (see page 29).

- 2.00 kg certified KIRIAL hybrid ryegrass
- 3.00 kg certified NOVIAL hybrid ryegrass
- 2.50 kg certified BOYNE perennial ryegrass
- 3.00 kg certified ABERBITE tet. perennial ryegrass
- 0.60 kg certified ABERDAI white clover
- 0.60 kg certified ABERSWAN white clover
- 0.30 kg certified ALICE white clover

12.00 kg/acre - £63.88

30.00 kg/ha - £159.70

## Additions



Red Clover: 1 kg red clover  
Cover Crop: 3 kg westerwold  
Heavy Land: 2 kg timothy  
Light Land: 2 kg cocksfoot  
Anti Bloat: 5 kg sainfoin

£8.60 per acre  
£8.55 per acre  
£10.70 per acre  
£10.30 per acre  
£14.00 per acre

## Milk-Meat Cut or Graze

## Five Year Plus Ley


Code: MIXMM

Our best selling dual purpose ley, equally suitable for cattle or sheep. This mixture combines the benefits of high yielding intermediate ryegrass Aberwolf, with Aberbite, which consistently holds its quality late into the season, along with highly palatable timothy and white clover. It can be sown for silage and hay or intensively grazed. This versatile high D-value ley will yield well on all soil types.

- 2.50 kg certified ABERWOLF perennial ryegrass
- 2.00 kg certified ABERBITE tet. perennial ryegrass
- 4.00 kg certified CALIBRA tet. perennial ryegrass
- 2.00 kg certified TODDINGTON perennial ryegrass
- 2.00 kg certified WINNETOU timothy
- 0.20 kg certified ABERDAI white clover
- 0.10 kg certified ABERSWAN white clover
- 0.20 kg certified ALICE white clover


13.00 kg/acre - £66.92

32.50 kg/ha - £167.30



Late Grazing  
Cornwall  
11th October





# Drought Resistant Leys

Drought can devastate forage crop yield. Avoid the severe consequences by choosing deep-rooting mixtures.

## Sowing and Growing

### Suitable soils and optimum pH

These mixes are designed for light, free-draining land with a pH of 5.6-7.

### When to sow

Sow between March and early September. Avoid late autumn sowing when mixtures contain clovers.

### How to sow

Sow into a fine, firm seedbed at around 10mm. Seed can be broadcast on a windless day, harrowed lightly and rolled. Alternatively, seed may be drilled in two directions into a well consolidated (rolled) seedbed.

### Management

These leys depend upon developing a large number of deep roots. To achieve this these leys should be allowed to accumulate a lot of leaf and should then be heavily grazed (rotationally) before being allowed to repeat the cycle. Set stocking is less effective. Leys containing cocksfoot should be grazed frequently and cut young to ensure that growth remains leafy.

### Nutrient requirements

N fertiliser (40kg N/ha) can increase early spring growth but high applications will impede clover growth and content which needs to be high for summer grazing production. P and K levels should be maintained at ADAS Index 2.

### Yield potential

Cocksfoot-based leys: 12t DM/ha

Ryegrass-based leys on dry, light land: 7t DM/ha

Ryegrass-based leys with rainfall: 12t DM/ha

The dry conditions suffered by many in recent years demonstrates the need for grass mixtures which continue to yield even during prolonged spells of drought. By combining deep-rooting grasses and clovers with differing growth habits, it is possible to provide summer grazing from dry soils.

Recent weather has shown that ryegrass does not thrive in dry conditions. However there are other grasses, such as meadow fescue, timothy and cocksfoot, which can be relied upon. These species can offer great benefits over ryegrass to those in challenging conditions. If you farm on dry land then these leys are well worth considering.

## Growing grass on droughty land

Pioneers of grassland farming, Robert H Elliot and William Lamin, developed complex ley mixtures comprising deep-rooting species such as cocksfoot, chicory and red clover.

Then, as now, some farmers were reluctant to use too much cocksfoot (see page 4), as it was inclined to grow coarse and clumpy. However, this is only a problem when seed is sown too thinly, allowing the cocksfoot too much freedom, or when it is allowed to become too mature when making hay.

Elliot observed first hand at Clifton Park that his deep-rooting four year ley provided good quality forage and improved soil so much that he was able to grow subsequent cash crops for four years with little fertiliser input. Lamin, who used a simplified version of Elliot's mix, observed "...it's like throwing money away to put ryegrass on dry land."

It's worth noting that although ryegrass is vulnerable to drought and is one of the first grasses to stop growing, it does recover very quickly when rain comes and can make a valuable contribution after prolonged dry periods.



## Mixes

## Light Land Beef/Sheep

## Long Term Ley

Code: MIX5

A long term ley that combines perennial ryegrass with drought-resistant cocksfoot. Good year-round production can be achieved which can be used for grazing or silage. Cocksfoot needs frequent grazing but recovers quickly after defoliation. The blend of persistent white clovers and timothy makes the sward extremely palatable. It is also very productive late into the summer.

- 3.00 kg certified DONATA cocksfoot
- 3.00 kg certified CALIBRA tet. perennial ryegrass
- 2.50 kg certified BOYNE perennial ryegrass
- 2.30 kg certified TODDINGTON perennial ryegrass
- 1.50 kg certified WINNETOU timothy
- 0.25 kg certified ABERSWAN white clover
- 0.25 kg certified ABERDAI white clover
- 0.20 kg certified ABERACE wild white clover

**13.00 kg/acre - £69.79**    32.50 kg/ha - £174.48

## Cholderton

## Four Year Plus Grazing/Cutting Ley

Code: MIXCM

A ley developed on the thin, chalk soils of Wiltshire which provides good growth for early grazing or cutting. It regrows powerfully through the spring and into the summer, giving an outstanding second cut yield. The ley tolerates dry conditions due to the deep roots of cocksfoot and red clover.

- 1.50 kg certified KIRIAL hybrid ryegrass
- 4.00 kg certified CALIBRA tet. perennial ryegrass
- 2.20 kg certified ABERWOLF perennial ryegrass
- 2.00 kg certified WINNETOU timothy
- 2.00 kg certified SPARTA cocksfoot
- 0.50 kg certified AVISTO red clover
- 0.40 kg certified ABERSWAN white clover
- 0.30 kg certified ABERDAI white clover
- 0.10 kg certified ABERACE wild white clover

**13.00 kg/acre - £68.53**    32.50 kg/ha - £171.33

## 'Lamins' Drought Resistant

## Four Year Grazing for Dry Land

Code: MIXCG04

This is a traditional humus building, drought resistant ley which is ideal for continuous grazing. This 'Clifton Park' type mixture will provide good quality forage which is high in protein. It starts early in the spring and will grow well through the summer and into the autumn. Birdsfoot trefoil has been added to this mixture after witnessing its ability to stay green throughout the dry summer of 2018.

- 5.50 kg certified DONATA cocksfoot
- 2.70 kg certified PARDUS meadow fescue
- 1.50 kg certified WINNETOU timothy
- 1.00 kg certified ALTASWEDE red clover
- 0.30 kg certified ABERSWAN white clover
- 0.30 kg certified ABERDAI white clover
- 0.20 kg certified LEO birdsfoot trefoil
- 0.50 kg certified PUNA II chicory
- 0.10 kg certified ENDURANCE ribgrass
- 0.25 kg burnet
- 0.05 kg yarrow
- 0.10 kg sheeps parsley

**12.50 kg/acre - £85.85**    31.25 kg/ha - £214.63

## Chicory Grazing Ley

## Three - Four Years

Code: MIXCL

This high-protein, mineral-rich, drought resistant mixture combines one of the most well-known varieties of chicory with clover and a small quantity of ryegrass. It will last for three to four years.

- 2.50 kg certified PUNA II chicory
- 0.20 kg certified ENDURANCE ribgrass
- 1.50 kg certified AVISTO red clover
- 0.60 kg certified ABERDAI white clover
- 1.70 kg certified HURRICANE tet. perennial ryegrass

**6.50 kg/acre - £71.00**    16.25 kg/ha - £177.50


## Additions



- Cover crop: 3 kg westerwolds
- Cover crop: 3 kg Italian ryegrass
- Cover crop: 10 kg vetches
- Anti bloat: 5 kg sainfoin

**£8.55 per acre**  
**£9.00 per acre**  
**£18.00 per acre**  
**£14.00 per acre**





# Herbal Grazing Leys

Deep rooting, species rich, nutritionally balanced grazing leys.

## What is a herbal ley?

A herbal ley is a complex seed mixture of grasses, legumes and herbs, which bring a range of benefits to forage, livestock health and soil fertility. Herbal leys can often include a mixture of up to 17 species, depending on the aims of the ley, location and soil type.

They have traditionally been used to build soil fertility and structure in an arable rotation, acting as a minimal input, four year break crop, but they bring significant benefits not only to the soil health, but also to the health and diet of livestock and the wider environment.

The deep rooting species in the mixture add drought tolerance when grown on thin soils or during dry summers, remaining green and palatable for much longer than other forage mixtures. They work especially well on dry, light land where ryegrass leys prone to burning up in mid summer.

The mixture of species also ensures a longer growing season and certain species included in the mixtures such as sainfoin, chicory and birdsfoot trefoil, have anthelmintic properties, which helps to reduce the worm burden in livestock, creating less reliance on artificial wormers.

The deep rooting herbs, notably chicory, mine the soil for important nutrients and minerals, making them available to the grazing livestock and lowering the need for bought in concentrates. The high legume content of these leys reduces the need for expensive artificial nitrogen, since they fix their own N, feeding the other grasses and herbs in the mixture, and again helping to reduce costs.

Newman Turner, one of the great advocates of herbal leys, described these mixes as his 'fertiliser merchant, food manufacturer and vet all in one'.

## Sowing and Growing

### Suitable soils and optimum pH

Ideally suited to medium and light soil types with a pH of 6.0-8.0.

### When to sow

Sow from March until early September.

### How to sow

Sow into a fine, firm seedbed after an application of FYM. These leys contain many small-seeded species and are best broadcast as this leads to more even plant distribution. Roll twice after sowing for maximum seed to soil contact and consolidation.

### Management


Graze lightly while the crop is establishing. Once growing well, rotationally graze allowing at least 28 days or more for recovery and regrowth. Using electric fencing, ration an area per day (e.g. about one acre for 100 cattle) but adjust this area to match growth and stock requirements. Over-grazing will damage chicory crowns. Surplus production from Herbal Leys can be made into silage.

### Nutrient requirements

No N is required, P and K should be maintained at ADAS Index 2.

### Yield potential

Yields of 13t DM/ha for the Herbal Ley should be achieved.



Interested in herbal leys? Learn more about their benefits and how they've stood the test of time in our 32 page farmers guide - **The Herbal Ley Farming System**

Visit [cotswoldseeds.com](http://cotswoldseeds.com) to download your copy.



## Mixes

Grass

Legume

Herb

## Simple Herbal Ley

## Four Year Grazing/Cutting/AD Ley

Code: MIX23

Our Simple Herbal Ley is designed for farmers who may be considering experimenting with a diverse seed mixture that's more complex than ryegrass and clover mixes. Our Simple Herbal Ley contains ryegrass, cocksfoot, timothy, meadow fescue and legumes, red clover, white clover, and mineral rich forage herbs including chicory and ribgrass.

- 2.00 kg certified LOFA festulolium
- 2.45 kg certified CALIBRA tet. perennial ryegrass
- 2.40 kg certified ABERWOLF perennial ryegrass
- 1.50 kg certified DONATA cocksfoot
- 1.50 kg certified WINNETOU timothy
- 0.80 kg certified PARDUS meadow fescue
- 0.30 kg certified ABERDAI white clover
- 0.20 kg certified ALICE white clover
- 0.25 kg certified AVISTO red clover
- 0.15 kg certified ERMO alsike clover
- 0.30 kg certified PUNA II chicory
- 0.15 kg certified ENDURANCE ribgrass

12.00 kg/acre - £68.60

30.00 kg/ha - £171.50

## Herbal Over-Seeding

## Deep-Rooting Herbal ley

Code: MIXHOS

Deep rooting herbal leys are becoming more and more popular. Grass-only swards lack protein rich clovers and mineral rich herbs. Ideally, herb-rich swards are best established by reseeding but where this is not possible this mixture can be oversown into a grass-only sward.

- 0.40 kg certified LEO birdsfoot trefoil
- 0.35 kg certified GLOBAL red clover
- 0.25 kg certified ERMO alsike clover
- 0.25 kg certified MERWI white clover
- 0.20 kg certified ABERDAI white clover
- 0.10 kg certified ABERACE wild white clover
- 0.20 kg commercial sweet clover
- 2.00 kg commercial sainfoin
- 1.00 kg burnet
- 0.50 kg sheeps parsley
- 0.35 kg certified LACERTA chicory
- 0.25 kg certified ENDURANCE ribgrass
- 0.15 kg yarrow

6.00 kg/acre - £54.28

15.00 kg/ha - £135.70

## Herbal Grazing Ley

## Four Year Drought Resistant Ley

Code: MIX20

Based on Newman Turner's original recommendations, this all round mixture provides wholesome and substantial forage for grazing and occasional cutting. It can provide grazing for early turnout and continues to produce forage right through the summer and autumn. Containing deep-rooting ingredients, this ley not only improves soil structure but also draws up essential vitamins and minerals for the ruminant animal.

- 1.60 kg certified DONATA cocksfoot
- 1.50 kg certified LOFA festulolium
- 1.00 kg certified ASTONKING perennial ryegrass
- 0.60 kg certified CALIBRA tet. perennial ryegrass
- 0.60 kg certified WINNETOU timothy
- 0.50 kg certified KORA tall fescue
- 0.40 kg certified PARDUS meadow fescue
- 2.50 kg commercial sainfoin
- 0.70 kg certified MILVUS red clover
- 0.60 kg commercial sweet clover
- 0.50 kg certified ABERDAI white clover
- 0.30 kg certified LUZELLE lucerne
- 0.20 kg certified ERMO alsike clover
- 0.20 kg certified LEO birdsfoot trefoil
- 0.70 kg burnet
- 0.60 kg certified PUNA II chicory
- 0.20 kg certified ENDURANCE ribgrass
- 0.20 kg sheeps parsley
- 0.10 kg yarrow

13.00 kg/acre - £87.25

32.50 kg/ha - £218.13

## Herbal Heavy Land Ley

## For Medium and Clay Soils

Code: MIX22

Still deep rooting but without cocksfoot this grazing mix suits heavier soils and lasts up to five years.

- 1.50 kg certified LOFA festulolium
- 1.70 kg certified NOVIAL hybrid ryegrass
- 2.20 kg certified CALIBRA tet. perennial ryegrass
- 1.40 kg certified PARDUS meadow fescue
- 2.25 kg certified WINNETOU timothy
- 0.80 kg certified KORA tall fescue
- 1.00 kg certified MILVUS red clover
- 0.65 kg commercial sweet clover
- 0.50 kg certified LUZELLE lucerne
- 0.50 kg certified ABERDAI white clover
- 0.40 kg certified ERMO alsike clover
- 0.45 kg certified PUNA II chicory
- 0.50 kg burnet
- 0.15 kg certified ENDURANCE ribgrass

14.00 kg/acre - £89.98

35.00 kg/ha - £224.95



Mob grazing herbal ley  
FarmED  
12th July



First Hand  
**Tom Day**



<b>Farm Type</b>	<b>Mixed</b>
<b>Location</b>	<b>Kent</b>
<b>Size</b>	<b>280 acres</b>
<b>Soil Type</b>	<b>Weald clay</b>
<b>Mixes Used</b>	<b>Herbal Ley, Maximum Yield Two Year Silage Ley, Forage Rape</b>

When the spring of 2020 brought drought, herbal leys were the only forage left for Tom Day’s sheep to graze.

Tom is a 4th Generation Farmer and his family has been farming the 280 acre tenanted farm in the Weald of Kent since 1900. Originally, the farm was mixed. ‘It had pretty much everything you could ever imagine being on a farm,’ says Tom. But gradually the less profitable enterprises got dropped and since Tom took over management in 2008 he has focussed predominantly on sheep as well as producing some hay and haylage for the local horse market. There are 450 breeding ewes; Romney and Romney crosses, with a Beltex tup to improve the carcass quality of lambs.

Weald clay is heavy and notoriously difficult to work. In the summer it bakes out, while in winter it’s wet, sticky and unworkable. Tom decided to try herbal leys after reading about them in the Cotswold Seeds catalogue.

‘I started using Cotswold Grass Seeds’ mixtures eight years ago because the catalogue is so informative, full of fresh ideas which helped me to think outside the box. I was keen to try herbal leys. They sounded great to repair and improve soil structure and workability and I liked the environmental benefits too.’

Tom has been using herbal leys for 3 years across 50 acres, so the farm now comprises 100 acres of permanent grass and pasture and the rest is in rotation between Cotswold Seeds’ maximum yield ryegrass for haylage and herbal leys, with forage rape as a break crop between the grass.

‘The sheep far prefer the leys to permanent pasture,’ Tom says. ‘We used to setstock and heavily graze a field then move them on, but now we’re learning to mob-graze. The clovers, legumes and grass comes back quicker and there is always a fresh bite in front of the sheep. It was also good to talk it all through with Cotswold Seeds who were always there at the end of the phone for advice and can tailor mixtures depending on what I need.’

Tom’s most recent herbal ley was sown last summer when weather conditions were right and the soil was most workable. It had established well by autumn and came back strong. The hot, dry spring of 2020 was especially difficult for reliable grass growth so Tom was glad he’d started the herbal leys journey.

**“A lot of the permanent grass just withered, shriveled and was gone. Thanks to the deep roots, the mob-grazed herbal ley was still going strong and providing plenty of food for the sheep.”**

‘A lot of the permanent grass just withered, shriveled and was gone. Thanks to the deep roots, the mob-grazed herbal ley was still going strong and providing plenty of food for the sheep.’

Tom says he definitely intends to carry on with herbal leys. ‘I like the variety of species and environmental benefits and the sheep do love it’





# Intensive Dairy

The latest ryegrass swards to maximise milk from grazed grass.

## Sowing and Growing

### Suitable soils

Ryegrass is a shallow-rooted grass and should be grown on moisture-retentive soils that do not dry out. The target soil pH for ryegrass grass leys is slightly acidic at around 6.

### When to sow

Ryegrass will germinate quickly from seed and sowings can be made from late February until early October in southern areas. However, most seed is sown in March and April and from late July until mid September when soil temperatures are above 7°C.

### How to sow

Drill in two directions into a fine, firm seedbed at 10-20mm. Rolling with either a Cambridge or flat roller before and after sowing is recommended. Broadcast seed should be harrowed lightly after sowing and before rolling.

### Management

Optimum sward height for grazing is 7-10cm. Early bite can be obtained on light, sheltered land from Intensive Dairy Graze - Early. Around 20 acres (8 ha) is advisable to provide early bite for 100 cows. Mid summer grazing is increased by grazing ryegrass hard early in the season as this stimulates tiller production and results in more leaves and less stem. Under-grazing should be avoided as this leads to stem production and loss of forage quality. To avoid winter damage, ryegrass should be grazed down to 4cm at the end of the season.

### Nutrient requirements

N fertiliser (70kg N/ha) will be required in mid February in mild areas to mid April for northern or upland districts. Provided soil moisture is available, a mid season N fertiliser application (60kg N/ha) will produce more grass. On land where late season grass can be utilised a further dressing of N (40kg N/ha) can be applied in late August. P and K should be maintained at ADAS index 2

### Yield potential

Yields of around 13t DM/ha are achievable if N levels are maintained and grazing is actively managed.

Grazed grass is by far the most important and economical feed and can provide around two thirds of forage on dairy farms.

Ryegrass mixtures provide very high annual yields when managed intensively in a high-input system. Ryegrasses allow for higher stocking rates than alternative grasses, such as meadow fescue or cocksfoot, because they are significantly more responsive to nitrogen fertiliser.

These mixtures include the latest varieties and grow very well on moisture-retentive 'ryegrass soils', providing grass from spring through summer and well into the autumn.

## Mixes

### Intensive Dairy Graze - Early

#### Four - Five Year Ley

Code: MIX3

To provide quality grazing throughout the season this mixture includes tetraploid ryegrasses which will increase palatability and voluntary intake. Containing early, mid and late season grasses, this ley is suited to light land, requires early grazing and can help to extend the grazing season. The inclusion of the 'Aber' strains ensures D-values remain high.

- 3.00 kg certified KIRIAL hybrid ryegrass
- 2.00 kg certified ABERWOLF perennial ryegrass
- 5.00 kg certified ABERCLYDE tet. perennial ryegrass
- 2.00 kg certified ABERBITE tet. perennial ryegrass
- 2.00 kg certified ASTONKING perennial ryegrass

**14.00 kg/acre - £64.90**    35.00 kg/ha - £162.25

### Intensive Dairy Graze - Late

#### Five Year Plus Ley

Code: MIX4

A proper late, thick-bottomed sward for summer grazing or late silage making on heavier soils. A high sowing rate of palatable grasses including high sugar 'Aber' varieties with consistent D values and late diploid DLF varieties with very good Crown Rust and Drechslera scores, to ensure grazing remains palatable late into the autumn. This mix costs more than other ryegrass mixes but provides an ideal grazing sward for at least five years.

- 2.00 kg certified ABERWOLF perennial ryegrass
- 2.00 kg certified BOYNE perennial ryegrass
- 4.00 kg certified ABERCLYDE tet. perennial ryegrass
- 4.00 kg certified ABERBITE tet. perennial ryegrass
- 3.00 kg certified TODDINGTON perennial ryegrass

**15.00 kg/acre - £74.95**    37.50 kg/ha - £187.38





# Long Term Leys

Good traditional leys that will produce grass for years to come.

Long term leys are ideal for self-sufficient beef and sheep farmers wanting to produce profitable stock with the emphasis on seasonal production, live weight gain and finishing healthy animals.

The long term ley mixtures contain perennial ryegrass or meadow fescue along with timothy and clover for good year-round production. Clover is an excellent protein source which increases production, reduces inputs and maintains profit margins.

All grasses have a lifespan. Some such as timothy and meadow fescue are very long lived and so can be considered permanent. Most ryegrasses are suited to short or medium term leys, but some varieties of late perennial ryegrass are persistent and suited to long term leys.

## Under Sowing a Spring Cereal

Since the advent of the Norfolk Four Course rotation, by Coke of Holkham, leys have often been sown underneath a spring sown cereal. This when done correctly protects the vulnerable new ley from hot weather and leaves a new ley well established after the cereal has been harvested. Any cereal can be under sown but barley and oats are most popular.

The competition factor should be reduced to a minimum by sowing the cereal, at two thirds the normal rate, and the ley seeds mix at the same time. The cereal should be drilled in and the grass seed mix broadcast on the surface, then harrowed and rolled. Some farmers wait until the cereal is up with 3 or 4 leaves before sowing as there is less risk of having a lot of green material going through the combine at harvest. This is really only advisable in high rainfall areas and not usually necessary in drier districts.

Sometimes, if the weather is bad, it is not possible to sow before the cereal has germinated. In this case sowing of the ley mix should be delayed until the cereal is through and well established with 3 or 4 leaves.

## Sowing and Growing

### Suitable soils and optimum pH

These leys are suitable for all soils apart from light ones. Ideally pH6-7.

### When to sow

These long-lasting leys take time to germinate and become established. It is therefore essential to sow when growing conditions are good and not too early in the spring before the soil is warm. They are often sown in spring but autumn sowings can be contemplated provided the seed is in by early September. These leys contain many small-seeded species and are best broadcast as this leads to more even plant distribution. Once sown, roll immediately to ensure good soil-to-seed contact.

### How to sow

With long term leys it is extremely important to control perennial weeds prior to sowing. As these mixes contain small seeded species such as timothy or clover it is best to sow at no more than 10mm into a fine seed bed. A cover crop of westerwolds ryegrass can provide additional bulk in the year of sowing but is not advisable on heavy ground as they may out-compete the other species in the mix. These mixes can alternatively be undersown to a spring cereal which has been drilled at a reduced rate.

### Management

Light grazing with sheep or young cattle will consolidate new plants, encourage grass to tiller and control annual weeds (known as the 'golden hoof'). Cutting for silage or hay is best left until the ley is well established in its second season.

### Nutrient requirements

These leys should receive no more than 100kg N/ha with the biggest demand (60kg N/ha) in March and April if a silage cut is to be obtained. Pochon Persistent requires very little or no N as the high clover content (30-50%) will fix N in the soil.

### Yield potential

Yields of 12t DM/ha should be achieved.



Grass

Legume

## Mixes

## Pochon Persistent

High Clover Long Term Grazing Ley Code: MIXCG01

For over thirty years Pochon has proven very successful on a wide range of conventional and organic farms. This mix is suitable for taking a cut of silage, but is mainly for rotational grazing. Including the best strains of high yielding white clovers from Aberystwyth, it gives excellent mid-summer production without artificial N.

- 3.00 kg certified TODDINGTON perennial ryegrass
- 2.00 kg certified ASTONKING perennial ryegrass
- 2.50 kg certified ABERBITE tet. perennial ryegrass
- 3.00 kg certified CALIBRA tet. perennial ryegrass
- 0.60 kg certified ABERSWAN white clover
- 0.60 kg certified ABERDAI white clover
- 0.30 kg certified ABERACE wild white clover

**12.00 kg/acre - £68.19** 30.00 kg/ha - £170.48

## Permanent Grass

Long Term Grazing or Cutting Code: MIX6

This versatile mixture contains persistent varieties ensuring good yields for cutting and grazing over many years. Timothy is extremely resistant to cold temperatures and provides good late-season growth. The thick-bottomed sward structure is obtained by using late perennial ryegrasses and highly nutritious white clover. This mixture is very hardy and can be grown in upland or lowland areas.

- 3.00 kg certified TODDINGTON perennial ryegrass
- 2.00 kg certified ASTONKING perennial ryegrass
- 2.50 kg certified ABERBITE tet. perennial ryegrass
- 2.50 kg certified ABERWOLF perennial ryegrass
- 2.00 kg certified WNNETOU timothy
- 0.40 kg certified ABERSWAN white clover
- 0.40 kg certified ABERDAI white clover
- 0.20 kg certified ABERACE wild white clover

**13.00 kg/acre - £69.90** 32.50 kg/ha - £174.75

## Long Lasting Upland

Dual Purpose Mix Code: MIXCG05

This ryegrass-free mix is very long lasting and will tolerate harsh upland conditions. It is very palatable and is best when rotationally grazed to allow a period of recovery and regrowth. It can also be cut for silage or hay with the best quality forage coming from swards which are cut before heading.

- 7.50 kg certified PARDUS meadow fescue
- 3.00 kg certified WINNETOU timothy
- 1.00 kg certified ALTASWEDE late red clover
- 0.40 kg certified ABERDAI white clover
- 0.40 kg certified ABERHERALD white clover
- 0.20 kg certified ABERACE wild white clover

**12.50 kg/acre - £79.35** 31.25 kg/ha - £198.38



Summer grazing  
Wiltshire  
30th June

## Additions




Westerwolds can provide cover during establishment and increase yields in the first year.

Add 3 kg of westerwolds

**£8.55 per acre**





# Root Crops

Fodder crops provide essential forage when grass is restricted. They are also a vital break crop.

Once sown, brassicas quickly produce a fodder crop. Adding muck or fertiliser to the crop make it as productive as possible. This then feeds a larger number of livestock, so returning more dung to the soil, making the most of a very beneficial cycle. Once the decision has been made to break up a ley or pasture, many farmers sow a brassica fodder crop. These are not troubled by grass pests or diseases and thrive on the nitrates released by the decaying sward.

## Reduce feed costs

These short-term catch crops are sown in late spring or summer to provide valuable home-grown fodder for buffer feeding dairy cows or finishing lambs in autumn or winter, when other sources of forage are limited. Turnips and rape grow quickly, needing just 10 weeks. Kale, swede and hardy turnip take a bit longer but are much more winter hardy and excellent for late-winter grazing. All are highly beneficial break crops which reduce grassland weeds and pest attacks.

## Summer feed for dairy cows

Stubble turnips are palatable, energy-rich and offer dairy farmers the opportunity to prevent a feed shortage over the summer. To allow the rumen to adjust, cows should be introduced gradually to the crop for the first few days.

## Lamb finishing

Lambs can be successfully fattened on fodder brassicas, gaining around 100-150 grams per day. The addition of a small quantity of hay, barley or concentrates is beneficial. Root crops, especially when grown on free-draining soils, are excellent for late autumn and winter use.

For more information on specific varieties visit [cotswoldseeds.com/knowledgehub.asp](http://cotswoldseeds.com/knowledgehub.asp)

## Sowing and Growing

### Suitable soils and optimum pH

These crops will grow on most soil types provided they are well-textured and can give a fine tilth when cultivated. However, it's important to sow on well-drained ground for winter grazing. Optimum pH6.2.

### When to sow

Fast growing root crops can be sown anytime from spring through till early autumn providing soil moisture is sufficient.

The slower growing crops such as the Hardy Root Mix, maincrop turnip, swede, fodder beet and kale should be sown in late spring (April – June).

### How to sow

Root crops (except fodder beet) can be direct drilled with a Moore Uni-Drill (or similar). If a root crop is to follow a grass ley, glyphosate can be applied to the grass prior to mowing. Once the grass is removed, seed can be direct drilled into its stubble and can establish quickly unhindered by weeds. A good dose of slurry or FYM should be applied before sowing if possible.

### Management

Electric fencing allows the crop to be fed at a controlled rate and should be long enough to give all stock access to the crop face. By doing this there is also less wastage through trampling. Ideally, a grass 'runback' should be provided for animals to lie on.

### Nutrient requirements

These crops need 70kg N, 50kg P and 50kg K per hectare.

### Yield potential

Species	DM/Ha	CP (%)	D-Value
Stubble turnip	4.5t	17	69
Maincrop turnip	6.0t	9	80
Swede	8.5t	11	82
Fodder beet	14.0t	12	78
Kale	9.0t	17	68
Forage rape	4.5t	19	65
Hybrid - Interval	5-8t	19	68
Hybrid - Redstart	6-8t	19	68



## Mixes

## Early Fold Root Mix

Fast Growing and Cheap Seed

Code: MIXEF

This is a fast growing mixture capable of producing up to 45 tonnes per hectare with a dry matter content of 10% in approximately 10-12 weeks. Three acres feeds 100 sheep (complete diet) or 50 cows (quarter of diet) for a month.

- 1.50 kg certified SAMSON stubble turnips
- 0.50 kg certified HOBSON forage rape

2.00 kg/acre - £7.58

5.00 kg/ha - £18.95

## Hardy Root Mix

Longer Term

Code: MIXHR

Combining different brassicas together in a mixture is often beneficial as the crop is more reliable and higher yielding. This longer term mixture needs 20 weeks or more to produce its full yield, but will provide winter hardy keep until February or March. Three acres feeds 150 sheep (complete diet) or 75 cows (quarter of diet) for a month.

- 0.65 kg certified PINFOLD kale
- 0.65 kg certified GREEN GLOBE hardy turnip
- 0.20 kg certified GOWRIE swede

1.50 kg/acre - £22.68

3.75 kg/ha - £56.70



## Straights

## Fodder Beet

This root crop provides a huge yield which is highly digestible and has a high energy content. It should be sown in April and, because the seed is pelleted, can only be sown with a precision drill. **Seed is only available in one acre packs (50,000 seed units).**

Robbos variety is suitable for leaf lifting harvester and feldherr for hand harvesting or grazing in situ.

Robbos £86.00 per acre

Feldherr £88.00 per acre

## Straights

Brassica

## Forage Rape

This protein rich green forage can be ready to graze in as little as 12 weeks and is ideal for fattening lambs. The Hobson variety is mildew resistant and front tolerant. Sowing rate 4.00 kg/acre.

Hobson (4.00 kg/acre)

£3.75 per kg

## Stubble Turnip

Turnips are grown in most areas of the UK as a highly digestible catch crop, ready within 10-12 weeks from sowing. Sowing rate 2.00 kg/acre.

Samson (Sheep grazing)

£3.80 per kg

Rondo (Cattle grazing)

£3.80 per kg

## Kale

Kale is high yielding, protein rich and winter hardy. Usually grazed between September and March, depending on sowing time. Full crop ready in 20 weeks. Sowing rate 2.00 kg/acre.

Pinfold (2.00 kg/acre)

£9.80 per kg

Maris Kestrel (2.00 kg/acre)

£16.50 per kg

## Hybrid Rape/Kale

Introduced to capitalise on the benefits of both rape and kale, this fodder crop is ready in 12 weeks from drilling. Many farmers favour this crop as it can offer good regrowth. Choose Redstart for winter hardiness and strong regrowth or Interval for good yields and high palatability. Sowing rate 3.00 kg/acre.

Redstart (3.00 kg/acre)

£8.00 per kg

Interval (3.00 kg/acre)

£4.60 per kg

## Maincrop Turnip

This is the hardier type of turnip which requires 20 weeks growth and is suitable for grazing late into the winter. Hardy turnips yield around a third more than stubble turnips. Sowing rate 2.00 kg/acre.

Green Globe (2.00 kg/acre)

£11.40 per kg

## Swede

This crop is ideally suited to cooler, wetter parts of the north and west of Britain. For stock or pot. Sowing rate 1.50 kg/acre.

Gowrie (1.50 kg/acre)

£44.50 per kg



First Hand  
**Geoff Sansome**



<b>Farm Type</b>	<b>Mixed</b>
<b>Location</b>	<b>Worcestershire</b>
<b>Size</b>	<b>100 acres</b>
<b>Soil Type</b>	<b>Light sandy loam and sandy silt loam</b>
<b>Mixes Used</b>	<b>Early fold root mix</b>

Hawford Farm in Worcestershire lies on the banks of the River Severn, where Geoff Sansome farms 100 acres on fairly light sandy loam and sandy silt loam soils, fattening about 60 head of cattle a year, some of which go out as boxed beef. There are also some cereals and a significant area of land is in higher tier environmental stewardship.

A 10 acre field is given over to the early fold stubble turnips and forage rape mix, following a crop of winter barley.

‘This is about the third year we’ve grown this crop,’ says Geoff. ‘We don’t have any sheep but we’ve got a very good neighbour who does. He’s an incredibly good shepherd with some cracking sheep. We were looking for a winter cover crop and he was looking for winter forage so it seemed like the perfect opportunity to put those two things together.’

The crop was established mid August and the shepherd will be bringing between 150-200 in lamb ewes onto it mid December and grazes through to March.

‘Management is incredibly simple from our perspective,’ says Geoff. ‘It was direct-drilled into stubble and we’ve never applied any fertiliser or insecticide. The only input this crop has had was a low rate of graminicide because there was a high level of barley volunteers due to this year’s harvest. So it’s a really low input crop. Payment is paid per head, per grazing day which more than covers all our costs and it gives us the benefit of what the sheep leave behind in terms of the following crop, which is the residual manurial value of the grazing and a very clean and even seed bed that we can direct drill into.’

The grazier has a quad bike mounted fencing system so it takes him just half an hour to fence out a block, marking out a week’s grazing each time. He checks them every day and so far hasn’t needed to provide any run back area.

‘Because of the light soil they seem to do fine just with the crop as it is,’ says Geoff. ‘They move across, taking all the leafy tops, then come back and take all the turnip roots in a second bite. Because of the free draining soil type they keep very clean without making a mess in the field or encountering foot problems. The grazier has been really happy with how they’ve performed over the last 2 years and last year he fattened some lambs quickly as well as grazing ewes.’

**“A great opportunity to provide stewardship for the soil and also provide useful grazing and cover all your costs.”**

After the sheep have finished, about mid March, they are quickly followed with a crop of spring barley, usually using a contractor with a Weaving Machinery direct drill.

‘We do sometimes get a bit of regrowth from little bits and pieces of stubble turnips, but the broad leaved weed herbicides in the spring barley controls that very easily.’

Geoff believes there’s a great opportunity for people who haven’t got livestock but want the associated soil benefits to bring them back onto the arable farm, helping to boost soil health.

‘With an increasing spotlight on what we do with our soils over the winter period in terms of green cover, runoff, erosion and nitrates etc. this is a great opportunity to provide stewardship for the soil, provide useful grazing and cover all your costs and a little bit more.’

It really is a win win for the grazier and the arable farmer.



# Walk & Talk at FarmED

For a great day out in the Cotswolds, visit the

## FarmED Programme of Events

[farm-ed.co.uk](http://farm-ed.co.uk)

Prices vary and some are free for Cotswold Seeds customers.



**FarmED**  
FARM & FOOD EDUCATION



Inspire

Inform

Share



# Equine

Horses' requirements are sometimes different to that of sheep and cattle. These mixes have been created specifically with equine needs in mind.

## Sowing and Growing

### Suitable soils and optimum pH

These mixes are suitable for most soil types, mixes containing forage herbs prefer lighter soils which are less grass dominant.

### When to sow

Sow from April to September when soil temperature is above 7°C. Ryegrass based mixes or surface mixes with high sowing rates can be sown later than non ryegrass grazing mixes.

### How to sow

Mixes can be broadcast or drilled but sown no deeper than 1cm. Always roll after sowing, if using a Cambridge roll, roll the field twice for maximum consolidation and soil contact.

Surface mix sowing rates vary. High rates are used on areas that receive lots of use and need fast establishment, low use areas or areas that are being patched up have scope to reduce the sowing rate.

### Management

New swards can be lightly grazed around five or six weeks after establishment. Overgrazing should be avoided, swards should be given regular rest periods of a few weeks throughout the growing season to recover.

Surface mixes can be topped 4-5 weeks after sowing and again 4 weeks later to encourage the sward to thicken and plants to tiller, creating a denser surface.

Annual weeds will disappear when grazed. Perennial weeds should be controlled prior to seeding. Selective herbicides can be used on docks, thistles & nettles.

### Nutrient requirements

Soil nutrient levels should be checked every 2 years and deficiencies corrected. If fertilising, take horses off for 2 weeks after application to allow the nutrients to be absorbed properly.

Slow release N fertiliser products are available to reduce a strong flush of grass directly after application e.g 'Paddock Royale' (approx 2x 25kg bags per acre, Yara Guidelines).



For over 40 years we have been supplying farmers and landowners with bespoke grass seed mixtures to serve many specific requirements, one of which is improving livestock health.

Pastures that contain a more natural diverse mix of plant species have lowered production costs through extending the grazing season and decreasing or, in some cases, eliminating the need for concentrate feed. Increasingly we are seeing similar needs within the equestrian industry. By taking a more holistic approach we can help horse owners and carers improve pasture quality and in turn improve their horse's health.

## Why is it so important to get your pasture right?

It is generally believed that wild horses had the ability to self-medicate, roaming freely to forage for different plant species that would naturally provide them with the macro and micro nutrients, minerals and vitamins that they need. Due to a number of factors such as herbicide use and popularity of aggressive species such as ryegrasses the diversity of species and consequently the nutritive value in swards has declined. It has become increasingly popular to address the shortfalls in the pasture by feeding concentrates and supplements, but this is not the healthiest option for the horse or for the environment.

There are a number of health issues that can be directly related to pasture management, nutritional content of the pasture and time spent eating grass or conserved grass (hay or haylage). These include laminitis, developmental orthopaedic disease, gastric ulcers, colic, respiratory diseases, mud fever, and wormer resistance. By getting the right mix of grass, legume and herb species for your land and your horse's needs the aim is for your horse to gain the majority, if not all, of its nutritional requirements from forage, relying less on concentrates and spending more time out in the field whatever the weather.

## Surface Mixtures

Good grass surfaces are key for exercising horses safely and effectively. High sowing rates create dense turfs which withstand heavy equine usage. The species chosen ensure a uniform surface and maximum cushioning for the horses without affecting performance, if managed correctly.

## Repair Mixtures

Over-seeding can be a useful, low cost way of regenerating existing pasture which has become thin and tired with age or damaged through over-grazing or use. The existing grass sward is a very competitive environment for any new seeds to establish so ryegrass is usually used for over-seeding because it is the quickest to germinate and suitably aggressive.



## Equine Grazing Mixes

### Standard Horse Pasture

#### Long-Term Grazing and Hay

Code: MIX7

This is our standard mix which provides a fast establishing dense turf for grazing or the occasional cut of hay. It does contain ryegrass so would not be suitable for horses or ponies that are prone to laminitis or Equine Metabolic Syndrome.

- 2.50 kg certified BOYNE perennial ryegrass
- 3.00 kg certified TODDINGTON perennial ryegrass
- 1.50 kg certified PARDUS meadow fescue
- 2.50 kg certified MELLORI creeping red fescue
- 2.00 kg certified WINNETOU timothy
- 1.50 kg certified EVORA smooth meadowgrass

**13.00 kg/acre - £68.00**    32.50 kg/ha - £170.00

### Natural Pony Paddock

#### Long Term with Herbs & No Ryegrass

Code: MIXPP

This non-ryegrass mix contains a very wide selection of grasses & herbs, improving drought tolerance, forage value and fibre content, providing a healthy, balanced diet. The absence of ryegrass species lowers the risk of laminitis due to a lower sugar and higher fibre content. This mix is slower to establish than those with ryegrass. Sow no later than mid September.

- 2.50 kg certified PARDUS meadow fescue
- 2.50 kg certified MELLORI creeping red fescue
- 1.85 kg certified WINNETOU timothy
- 1.75 kg certified EVORA smooth meadowgrass
- 1.50 kg certified KORA tall fescue
- 1.50 kg certified SPARTA cocksfoot
- 1.00 kg certified BORNITO sheeps fescue
- 0.25 kg certified HIGHLAND common bentgrass
- 0.25 kg certified TENO smaller catstail
- 0.25 kg certified DASAS rough stalked meadowgrass
- 0.10 kg commercial meadow foxtail
- 0.05 kg commercial sweet vernal grass
- 0.10 kg certified ENDURANCE ribgrass
- 0.25 kg burnet
- 0.05 kg yarrow
- 0.10 kg sheeps parsley

**14.00 kg/acre - £89.80**    35.00 kg/ha - £224.50

### Racecourse, Gallop and Cross Country

#### Permanent

Code: MIX8

This mixture is very dense and resilient and contains spreading grasses with the ability to fill bare patches.

- 25% certified PLATINUM dwarf perennial ryegrass
- 25% certified EVORA smooth meadowgrass
- 50% certified MELLORI creeping red fescue

**50-160 kg/acre 125-400 kg/ha**    **£4.79 per kg**

### Equine Pasture Mix

#### Long Term and No Ryegrass

Code: MIX13

This is a persistent, non-ryegrass mix providing good quality forage with a low sugar content. Although this mix will take a little longer to establish than a ryegrass based mix, it will provide a dense, resilient turf with balanced forage for grazing and hay.

- 3.65 kg certified PARDUS meadow fescue
- 2.65 kg certified WINNETOU timothy
- 2.60 kg certified EVORA smooth meadowgrass
- 2.50 kg certified MELLORI creeping red fescue
- 1.40 kg certified KORA tall fescue
- 0.50 kg certified HIGHLAND common bentgrass
- 0.50 kg certified DASAS rough meadowgrass
- 0.20 kg certified TENO smaller catstail

**14.00 kg/acre - £83.69**    35.00 kg/ha - £209.23

### Paddock and Gateway Repair

#### Over-Seeding

Code: MIXPAD

A high sowing rate of fast-establishing mixture which provides a thick grass cover for use on poached areas or to improve pasture quality.

- 4.00 kg certified JIVET westerwold ryegrass
- 4.40 kg certified TODDINGTON perennial ryegrass
- 4.30 kg certified PLATINUM dwarf perennial ryegrass
- 4.30 kg certified MELLORI creeping red fescue
- 3.00 kg certified EVORA smooth meadowgrass

**20.00 kg/acre - £90.69**    50.00 kg/ha - £226.73

## Over-seeding Mixes

### Pasture Over-Seeding

#### Longer Term Four to Five Years

Code: MIXOSH

A long duration ryegrass mixture for over-seeding grazing pastures where more forage is required.

- 5.00 kg certified ASTONKING perennial ryegrass
- 5.00 kg certified TODDINGTON perennial ryegrass

**10.00 kg/acre - £52.50**    25.00 kg/ha - £131.25

### Equine Over-Seeding

#### Longer Term Four to Five Years

Code: MIXEQOS

A long duration non-ryegrass mixture for over-seeding grazing pastures where more forage is required.

- 4.80 kg certified PARDUS meadow fescue
- 2.20 kg certified WINNETOU timothy
- 2.00 kg certified MELLORI creeping red fescue
- 1.00 kg certified EVORA smooth meadowgrass

**10.00 kg/acre - £54.59**    25.00 kg/ha - £136.48



# Cover Crops

## Protecting and enhancing our soils

Rising cost of fertilisers and other inputs combined with an increasingly uncertain climate means that it's becoming ever more beneficial and indeed crucial to have good soils which are fertile, well structured and tolerant to drought.

The dilemma for many arable farmers is affording the time and money to invest in soil health and improvement, but simply put, if we don't, soils will rapidly become so poor they will adversely affect yield, crop quality and ultimately profitability.

It is the small remains of plant life that ultimately produce vital fertility and structure in the soil. Though this organic matter makes up only a small percentage of the soil content it is vital, since nothing will grow on clay, silt or sand alone. Doing nothing to replace it is therefore not an option.

The best and cheapest way of adding organic matter to the soil is to grow cover crops between cash crops. The cost of the seed and having no income from a field given over to cover crops can appear prohibitive but should be seen as an investment that will produce higher and better yields in subsequent years.

## Sowing and Growing

### Suitable soils and optimum pH

These will grow on most soil types with a pH above 5.6.

## How to sow


Mixes can be broadcast or drilled at a maximum of 10mm. Ideally, into warm soils when rain is imminent. If possible, roll after sowing for maximum seed to soil contact.

If sowing multi species mixes with large and small seeds drill according to the smallest seed size, sowing these too deeply will reduce establishment reliability.

Catch crops must be established by 20th August and maintained for a minimum of 8 weeks and retained until at least the 14th October. Cover crops must be established by 1st October to 15th January or longer.

These mixtures can be grazed, topped or sprayed off after the above dates to terminate and either incorporated or left to break down on the surface depending on the farming system.

Cover crops grown after high yielding cereals may benefit from 25-50kgs per ha of N, especially if sowing for winter grazing.



Discover a whole lot more on cover crops and green manures in our updated practical guide - **Sort Out Your Soil 2nd Edition**

Call us on 01608 652552 or visit  
cotswoldseeds.com to get your copy.



Legume

Herb

Brassica

Cereal

Other

## Short term mixes

## Quick Growth Cover Crop

Flexible, Fast Establishment

Code: MIXQCC

A mixture designed to establish rapidly, providing a flexible option to sow between an early harvested crop but before autumn planting, or as an autumn cover crop for rapid green winter cover.

- 2.45 kg certified ASTA mustard
- 2.30 kg certified IKARUS fodder radish
- 0.25 kg certified NATRA phacelia

5.00 kg/acre - £16.96

12.50 kg/ha - £42.40

## Diverse Grazable Cover Crop

Ground Cover &amp; Livestock Forage

Code: MIXGCC

A great mix for a joint arable & livestock venture, the mix contains soil conditioning phacelia, fodder & tillage radish, combined with high protein vetch, forage rape & turnips improving palatability and liveweight gain for stock. N.B. Rye will be replaced with winter cereal if unavailable early in the season.

- 7.30 kg certified PROTECTOR rye
- 3.50 kg certified JOSE common vetch
- 0.05 kg certified HEUSERS OSTSAAT crimson clover
- 0.01 kg commercial balansa clover
- 0.50 kg certified IKARUS fodder radish
- 0.30 kg certified HOBSON forage rape
- 0.15 kg certified SAMSON stubble turnip
- 0.15 kg certified RONDO stubble turnip
- 0.04 kg certified NATRA phacelia

12.00 kg/acre - £18.30

30.00 kg/ha - £45.75



Spring sown grazable cover crop  
Wiltshire  
1st August





# Green Manures

Protecting and enhancing our soils.

## Summer Sown Mixes

Summer green manures are planted from late spring onwards on bare ground and are incorporated before the sowing of a winter cash crop. A good summer green manure will be ready for turning-in after only 8-10 weeks. These crops give good leaf canopy cover to block out light, suppressing weed growth.

These green manures can be sown on their own or as an understory to a main crop and last between 2 and 6 months. As legumes will only fix nitrogen when the soil is above 8°C they are effective between April and August.

## Overwinter Mixes

Winter green manures such as rye or westerwolds scavenge excess nitrogen from previous crops which prevents it leaching over the winter. The nitrogen held within the green manure crop is then released when it is incorporated.

Legumes like vetch can be used for winter cover and, provided that these are sown by September, can fix up to 200kg N/ha for use by the following cash crop. The canopies of these plants also protect against soil erosion.

## Longer Term Leys

Slower growing perennial legumes such as red and white clover are used to add nitrogen to the soil over a long period.

Red clover fixes upwards of 200kg N/ha which is released rapidly after incorporation. To delay the release of nitrogen, clover is mixed with grass which is higher in carbon and acts like a sponge, holding the nitrogen for longer. This is especially important for subsequent autumn-sown crops where the nitrogen demand is highest 6 or 7 months after the green manure crop.



Summer green manure  
Worcestershire  
15th August

## Sowing and Growing

### Suitable soils and optimum pH

These will grow on most soil types with a pH above 5.6.

### When to sow

Sow summer mixes in warm soil between May and July. If undersowing, seed should be broadcast from mid March in damp conditions before the host crop canopy closes in. Cover for the winter should be sown by late September although rye and vetch can be sown into October.

### How to sow

Rye and vetch seeds can be drilled at up to 25mm. All other mixes should be drilled or broadcast at no more than 10mm.

### Management

Summer green manures will be ready for incorporation after 8-10 weeks normally at the onset of flowering. Winter green manures can be incorporated in April or May. Westerwolds ryegrass will regrow after cutting so can be left through the summer for further cutting or mulching. To minimise the risk of ryegrass seed being shed, cut before the seed heads are visible.

### Yield potential

The amount of N fixed by legumes depends on the success of the green manure. Generally, a reasonable crop can fix over of 100kg N/ha from a spring or summer sowing. Rye can scavenge and hold 90% of soil N, westerwolds about 70% and vetch and red clover can fix upwards of 200kg N/ha if left to grow.



Brassica

Legume

Herb

Cereal

Grass

## Short term mixes

## Summer Quick Fix

## Nitrogen Boost

Code: MIXSQF

The purpose of this mixture is to build soil N in a short time. It is a fast-growing, annual mixture that is at its best when sown into warm soils.

- 1.80 kg certified ASTA mustard
- 1.50 kg certified HEUSERS OSTSAAT crimson clover
- 1.20 kg certified AKENATON berseem clover
- 0.65 kg commercial sweet clover
- 0.60 kg certified LASER persian clover
- 0.25 kg certified GLOBAL red clover

6.00 kg/acre - £29.10

15.00 kg/ha - £72.75

## Summer Green Manure

## Early Sown N-Fixer

Code: MIXCCE

In warm soils, this mix can provide 150kg of N per hectare from a summer sowing. Very rapid growth with the potential to leave in over winter.

- 1.25 kg certified SHAKIRA Italian ryegrass
- 0.85 kg certified HEUSERS OSTSAAT crimson clover
- 0.65 kg commercial sweet clover
- 0.45 kg certified AKENATON berseem clover
- 0.10 kg certified GLOBAL red clover
- 0.90 kg certified ASTA mustard
- 0.65 kg certified IKARUS fodder radish
- 0.15 kg certified STRUCTURATOR tillage radish

5.00 kg/acre - £21.80

12.50 kg/ha - £54.50

## Yellow Trefoil/White Clover

## Intercrop Mixture

Code: MIXINT

This mixture will fill the base of a main crop brassica or cereal without affecting its yield. It reduces weed competition, adds organic matter and fixes nitrogen. Trefoil rarely interferes with harvest as it is low growing. This strong growth can eliminate weeds, especially if left in for a second year.

- 1.05 kg certified VIRGO PAJBERG yellow trefoil
- 1.95 kg certified MERWI white clover

3.00 kg/acre - £35.35

7.50 kg/ha - £88.38

## Over winter mixes

## Rye/Vetch

## Overwinter Mix

Code: MIXRYEV

Growing a N lifter and fixer together is a reliable way of improving soils over the winter. An excellent weed suppressor. Available from September.

- 55.00 kg certified ELEGO rye
- 20.00 kg certified JOSE vetch

75.00 kg/acre - £89.90

187.50 kg/ha - £224.75

## Ryegrass/Vetch

## Overwinter Mix

Code: MIXWWW

An economical, effective option for overwinter soil management. Adds large amounts of N and organic matter.

- 8.00 kg certified JIVET westerwolds ryegrass
- 17.00 kg certified EARLY ENGLISH vetch

25.00 kg/acre - £53.40

62.50 kg/ha - £133.50

## Longer term mixes

## Fertility Builder

## One-Two Year Mixture

Code: MIXFB

A grass and clover mix is the most effective green manure of all for improving soil fertility and structure. To realise its full potential it should be grown for at least one full year before incorporation.

- 2.65 kg certified GLOBAL red clover
- 0.50 kg certified MERWI white clover
- 5.85 kg certified CALIBRA tet. perennial ryegrass

9.00 kg/acre - £53.40

22.50 kg/ha - £133.50

## Humus Builder

## 2-4 Year Soil Structure Improver

Code: MIXHB

This mix utilises species with very strong tap roots for huge improvements to soil structure and organic matter levels, ideal on light or dry land.

- 4.00 kg certified GLOBAL red clover
- 0.50 kg certified LACERTA chicory
- 3.00 kg certified SPARTA cocksfoot

7.50 kg/acre - £57.73

18.75 kg/ha - £144.33



First Hand

## Mike Fisher



<b>Farm Type</b>	Market Garden
<b>Location</b>	Hampshire
<b>Size</b>	5 Hectares
<b>Soil Type</b>	Light loam over chalk
<b>Mixes Used</b>	Rye & Vetch, Winter Cover Crop, Herbal Leys & Bespoke Clover

Veg grower, Mike Fisher, has not applied any compost for the past five years, relying instead purely on plant mixtures as green manures.

This year, 2021, Mike celebrates 30 years as a grower. He began with a bare field and launched a veg box scheme back in 1994, one of the first organic growers. For a time he sold wholesale to Able & Cole and Riverford but now concentrates on the local market, harvesting once a week and supplying 130 boxes a week direct to households within a 10 mile radius. There's a variety of veg, including sweetcorn, leeks, plus tomatoes, cucumbers and salad leaves in polytunnels.

Unusual in his choice of growing organically, Mike was also a pioneer in the use of green manures and turned to Cotswold Seeds in 1992 because in those days they were the only seed merchant offering these mixtures.

'The first crop I used was forage rye to increase organic matter and soil life,' Mike remembers.

Now Mike uses a range of green manures in rotation, with 4 years of vegetables and 1 year of forage rye and vetch and red & crimson clover for building organic matter

and adding diversity. It starts with potatoes and these come out in September. Then garlic and winter onions go in until July. The red and crimson clover goes in before August and comes out in March before brassicas.

'It's a case of organising the rotation to provide a block of land for green manures,' Mike says. 'There are early and late planting and different harvest dates for veg so you can play about with the timing and the veg successions.'

He's experimented with buckwheat to control couch grass for the Organic Research Centre, which had some success. This year he sowed later so doubled up with phacelia and at the time of writing has yet to see if it smothered the couch. He's also used a Cotswold Seeds Herbal Ley. 'It might seem odd for a veg grower to plant herbs but it's important to bring up minerals and trace elements from the soil.'

Mike also adapts seed rates. 'We tend to use double seed rates to get good coverage and smother all our annual veg weeds. Having been growing intensive veg for 30 years it can be an issue, so worth the extra cost.'

**"The green manures not only build soil organic matter but also improve water retention too. For summer veg crops that is vital."**

The 2020 winter cover crop of radish, mustard, vetch, crimson clover went in the ground on 20th August. The crop was sown straight after two thunderstorms when the soil temperatures were high. A couple more thunderstorms made conditions ideal for establishment. The land had a ley on it recently, so some fertility was carried over from that and just 2 months later the cover crop was vast, reaching waist height.

'It's gone in ahead of next year's leeks,' Mike says. 'It will come out late February or March, flailed off then ploughed or rotavated.'

The results of Mike's rotation are impressive. The soil drains well and has a high pH of 7.8 and organic matter is now 9%. 'It was 5% immediately after stubble, so we weren't starting from nothing, but the improvement is very impressive.' It's up to 20% in the ½ acre of undercover polytunnels.

The green manures not only build soil organic matter but help with water retention too. 'For summer veg crops that is very important as well,' Mike says.

The proof is in the products. Many customers have been with Mike since the '90s and this year he has been inundated with new inquiries. 'I like building loyalty,' he says.


He's also a loyal customer of Cotswold Seeds. 'I have stayed with them for decades because of the great service and the quality of information and technical advice.'



# Environmental

## Countryside Stewardship, Entry Level & Higher Level Seed Mixtures.

Environmental seed mixtures are one way of protecting and enhancing wildlife across farmland. Many existing entry level & higher level stewardship schemes are still providing important resources and habitats. The more recent Countryside Stewardship scheme offers a further range of options, some based on the ELS/HLS prescriptions as well as newer options like the Bumblebird mixture and Two Year Legume Fallow. The mixtures below are common environmental stewardship prescriptions for ELS, HLS and Countryside Stewardship. All options can be tailored to better suit the location, soil type and aim of the scheme.



Interested in herbal leys? Learn more about their benefits and how they've stood the test of time in our 32 page farmers guide - **The Herbal Ley Farming System**

Call us on 01608 652552 or visit [cotswoldseeds.com](http://cotswoldseeds.com) to get your copy.

Sowing & Growing guide for **ALL** environmental mixtures follows on page 44.

### Stewardship scheme codes

CSS - Countryside Stewardship Scheme

ELS - Entry Level Scheme

HLS - Higher Level Scheme

### Mixes

#### GS4 Legume & Herb Rich Sward

ELS/HLS/CSS Codes: GS4, EK21

##### Whole Field Herbal Ley

Code: MIXGS4

This all round mixture promotes biodiversity, creates habitats, produces pollen and nectar and is also a superb soil conditioner and top quality forage. It contains deep rooting species which are drought resistant and also draws up micronutrients from deep within the soil.

- 12% certified cocksfoot
- 12.70% certified tet. perennial ryegrass
- 5.50% certified perennial ryegrass
- 10% certified timothy
- 5% certified meadow fescue
- 5% certified tall fescue
- 2.50% certified alsike clover
- 3% certified red clover
- 2% certified late flowering red clover
- 2% commercial sweet clover
- 20% commercial sainfoin
- 3% certified birdsfoot trefoil
- 4.50% chicory
- 6.50% burnet
- 1% yarrow
- 4% sheeps parsley
- 1.20% ribgrass
- 0.1% lesser knapweed

10.00 kg/acre 25.00 kg/ha

£6.88 per kg



Legume & herb rich sward  
Gloucestershire  
15th July



# Pollen & Nectar

Legume and flower margins.

## Margin mixes

### Flower Rich Margin AB8

ELS/HLS/CSS Codes: EC24, EE1, EE2, EE3, EE9, EF1, HE10, AB8

#### Long Term Pollen & Nectar

Code: MIXAB8

This straightforward, low cost mixture is ideal for awkward areas in arable fields. Relatively slow to establish, after the first year there are flowers for insects, seeds for birds and cover for mammals. This mixture meets the criteria for AB8 requirements pre-2020.

- 5% certified common bentgrass
- 10% certified smaller catstail
- 25% certified smooth stalked meadowgrass
- 25% certified slender creeping red fescue
- 30% certified sheeps fescue
- 1% ox-eye daisy
- 1% wild carrot
- 1% lesser knapweed
- 0.5% red campion
- 0.5% musk mallow
- 0.4% ribwort plantain
- 0.4% yarrow
- 0.2% self heal

**8.00 kg/acre** 20.00 kg/ha **£14.20 per kg**

### Pollen & Nectar Margin

ELS/HLS Codes: EE1, EE2, EE3, EE9

#### Grass and Legumes

Code: MIXPN

The mixture below is designed for ELS Pollen & Nectar margins but can also be used for HLS. It is best on light soil and lasts for around four years. Excellent for bumblebees and butterflies.

- 5% certified common bentgrass
- 5% certified crested dogstail
- 15% certified sheeps fescue
- 15% certified smooth stalked meadow grass
- 20% certified red fescue
- 20% certified meadow fescue
- 5% certified late flowering red clover
- 5% commercial sainfoin
- 4% certified common vetch
- 2% certified alsike clover
- 2% certified birdsfoot trefoil
- 2% certified yellow trefoil

**8.00 kg/acre** 20.00 kg/ha **£6.65 per kg**

### Floristically Enhanced AB8 Field Margin

ELS/HLS/CSS Codes: EF1, EE12, AB8

#### Long Term Pollen & Nectar

Code: MIXAB820

This mixture is fully compliant with the new 2021 scheme requirements for AB8, containing the required 4 grass species and 10 wildflower species. This mixture will provide a rich pollen and nectar source for pollinating insects such as bees, and butterflies throughout the summer months.

- 5% certified common bentgrass
- 5% certified smaller catstail
- 10% certified slender creeping red fescue
- 20% certified smooth meadowgrass
- 25% certified red fescue
- 25% certified sheeps fescue
- 1.85% salad burnet
- 1.50% ox-eye daisy
- 1.30% lesser knapweed
- 1.15% wild carrot
- 1% native red clover
- 1% red campion
- 0.90% white campion
- 0.75% ribwort plantain
- 0.20% yarrow
- 0.20% self heal
- 0.15% musk mallow

**8.00 kg/acre** 20.00 kg/ha **£17.96 per kg**

### The Operation Pollinator

ELS/HLS/CSS Codes: EF4, AB1

#### Just Legumes

Code: MIXOP

Operation Pollinator is a mixture of legumes without grasses to provide a flower-rich area. This mix works well on heavy soil types where sown grasses can become dominant.

- 26% certified late flowering red clover
- 22% certified red clover
- 20% certified alsike clover
- 20% commercial sainfoin
- 7% certified birdsfoot trefoil
- 4% certified crimson clover
- 0.35% wild carrot
- 0.35% ox-eye daisy
- 0.30% lesser knapweed

**5.00 kg/acre** 12.50 kg/ha **£8.99 per kg**



Grass

Legume

Herb

Cereal

Wildflower

Others

Brassica

# Farmland Birds

Reducing the hungry gap.

## Farmland bird mixes

### AB15 Autumn Sown Two Year Legume Fallow

ELS/HLS/CSS Codes: AB15

#### Two Year Mixture

Code: MIXAB15

Flowering crops on fallow land. Substitute perennial ryegrass for cocksfoot on dry soils.

- 52% tet. perennial ryegrass
- 14% perennial ryegrass
- 8% late flowering red clover
- 6% red clover
- 6% birdsfoot trefoil
- 13.7% common vetch
- 0.30% lesser knapweed

12.00 kg/acre 30.00 kg/ha

£5.76 per kg

### AB9 One Year Winter Bird Food

ELS/HLS/CSS Codes: AB9, EF2

#### Survival Mixture

Code: MIXSM

This mixture should be sited on field margins or next to hedges or woodland. It contains a range of species which provides food for birds, including finches and sparrows, over one winter while also feeding small mammals.

- 16% quinoa
- 16% fodder radish
- 14% white millet
- 14% red millet
- 14% linseed
- 10% mustard
- 10% spring barley
- 6% gold of pleasure

5.00 kg/acre 12.50 kg/ha

£4.76 per kg

### AB9 Two Year Wild Bird Seed

ELS/HLS Codes: AB9, EF2, CSS

#### Farmland Bird Feeder

Code: MIXWBSS

This mix includes cereal and quinoa for the first winter, while kale provides late seed in the second winter. It is a good food source for wild birds and is reliable if managed properly, as well as being more economical.

- 70% spring triticale (supplied separately)
- 14% game kale
- 10% white millet
- 2% quinoa
- 2% fodder radish
- 2% mustard

20.00 kg/acre 50.00 kg/ha

£3.04 per kg

### AB15 Autumn Sown Fallow Legume Only Mix

ELS/HLS/CSS Codes: AB15

#### Two Year Mixture

Code: MIXAB15LEG

Legumes only on fallow land. Use this if grass weeds are an issue.

- 50% common vetch
- 15% certified lucerne
- 10% alsike clover
- 10% late flowering red clover
- 6% red clover
- 5% birdsfoot trefoil
- 4% yellow trefoil

8.00 kg/acre 20.00 kg/ha

£5.66 per kg

### AB16 Enhanced Autumn Sown Bumblebird

ELS/HLS/CSS Codes: AB16

#### Two Year Mixture

Code: MIXAB162021

Food source for birds, pollinators and insects.

- 35% rye
- 18.75% winter wheat
- 15% winter barley
- 7.5% common vetch
- 6.5% winter linseed
- 2% crimson clover
- 2% fodder radish
- 4% game kale
- 1.5% lucerne
- 1.25% fennel
- 1.25% birdsfoot trefoil
- 1.25% gold of pleasure
- 1.25% alsike clover
- 0.75% phacelia
- 0.75% balansa clover
- 0.75% red clover
- 0.5% late flowering red clover



20.00 kg/acre 50.00 kg/ha

£2.81 per kg

### Retrieve Mix

ELS/HLS Codes: EC24, EE1, EE2, EE3, EE9, EF1, HE10, CSS

#### Fast and Economical

Code: MIXRET

For a summer sowing after a failed spring crop nothing beats these fast growing brassica species. It's quick, reliable and it works.

- 48% mustard
- 26% forage rape
- 16% fodder radish
- 8% hybrid rape/kale
- 2% brassica carinata

6.00 kg/acre 15.00 kg/ha

£3.32 per kg



# Resource Protection

Grassy areas to shield water courses and provide wildlife habitats.

## Sowing and Growing - Environmental Mixes

### Suitable Soils

Heavy, wetter soils can cause a dominance of strong grasses, consider the grass free Operation Pollinator where this is an issue, to maximise flowering species.

### When to Sow

For grass only or pollen & nectar mixes sow between late March and early May, or August and early September. Legume based mixtures should be sown into warm soils. Generally mixes containing wild flowers are sown in the autumn, especially if they contain Yellow Rattle.

Sow Wild Bird seed mixtures when the risk of frost has passed in the spring, fast growing patch up mixtures for failed spring sowings can be sown in August.

### How to sow

Species included in pollen and nectar or grass mixtures are small in size and should be shallow sown into a fine but firm seedbed. They can be broadcast and harrowed or shallow drilled at 10mm. Both options should be well rolled after sowing for maximum seed to soil contact.

Wild Bird seed mixtures containing a range of annual species can be shallow drilled or broadcast and well rolled in to a fine but firm seedbed.

Two year Wild Bird mixtures containing high levels of cereals can be supplied separately, the cereals can be drilled in rows to a depth of 25mm, and the smaller seeds broadcast in a 2nd pass.

### Management

Pollen & nectar and grass only mixtures can be lightly topped several times during establishment, normally 6-8 weeks after sowing, to control annual weeds and encourage tillering. They are also cut back as growth slows in the autumn.

Weed control in Wild Bird mixtures is difficult, it's important to create a clean, weed free seedbed before sowing. Once established the annual species will be left to grow through the winter, before being replanted the following year. Mixtures including Kale can be left for a 2nd year.

Stewardship agreement permitting, 50-60kgs/ha of fertiliser can boost seed yields and speed up establishment, ensuring leafy brassica crops grow through the risk of flea beetle damage. Some agreements allow herbicide control for weeds.

## Mixes

### Species Rich Parkland Grassland

ELS/HLS Codes: HK7, GS7

#### Low Maintenance Long Term

Code: MIXPGLM

A slow growing and manageable seed mix for those with low requirements from permanent grassland. This mixture can be grazed periodically or topped to keep a tidy appearance.

- 3% commercial meadow foxtail
- 5% commercial crested dogstail
- 1% commercial sweet vernal grass
- 5% certified common bentgrass
- 10% certified smaller catstail
- 22% certified smooth meadowgrass
- 24% certified red fescue
- 30% certified sheeps fescue

**16.00 kg/acre** 40.00 kg/ha **£8.62 per kg**

### Recreating Grassland

ELS/HLS Codes: HK7, ED2

#### Long Term

Code: MIXRG

The mixture below is suitable for sowing on most soil types ranging from clays to calcareous. Provides grass for grazing or hay production (if no forage is required use the mix above).

- 5% certified common bentgrass
- 10% certified sheeps fescue
- 15% certified red fescue
- 20% certified smooth meadowgrass
- 20% certified timothy
- 30% certified meadow fescue

**10.00 kg/acre** 25.00 kg/ha **£5.85 per kg**

### Buffer Strip Grass Margin

ELS/HLS/CSS Codes: EJ5, EJ9, EE7, EE8, EC24, EE1, EE2, EE3, EE9, AB3, SW4, SW1

#### Two, Four or Six Plus Metre

Code: MIXGM

An ideal mixture for buffer strips on cultivated land. This mixture is suitable for ELS and can also be used in HLS.

- 5% certified common bent
- 10% certified cocksfoot
- 20% certified timothy
- 20% certified meadow fescue
- 20% certified smooth meadowgrass
- 25% certified red fescue

**10.00 kg/acre** 25.00 kg/ha **£5.82 per kg**



First Hand

**David Gow**

<b>Farm Type</b>	<b>Mixed</b>
<b>Location</b>	<b>Oxfordshire</b>
<b>Size</b>	<b>1000 acres</b>
<b>Soil Type</b>	<b>Light sand to heavy clay</b>
<b>Mixes Used</b>	<b>Wild flowers</b>

David Gow has truly embraced the concept of farm diversification, branching out into selling Christmas trees and planting willow trees for cricket bats, which can only be grown successfully in the UK. But environmental concerns drive all the different enterprises.

David's 1000 acre farm is 5 miles west of Oxford and his family have been there since 1880, as tenants on some of the land and owning the rest. Two kilometres of the farm is bordered by the River Thames. There are 300 acres of ancient woodland, with 120 acres SSI and the rest is given over to a variety of crop. Six hundred acres of arable land grows winter wheat and maize as a break crop. There's a herd of beef suckler cows, a camping and caravan site and farm shop which sells the Christmas trees the Gow family have been growing since the 1950s.

Five years ago, David further diversified with a different cash crop, introducing willows trees for cricket bats in an irrigated set aside field. Willow trees are planted with a 10m spacing so there's bare ground between the trees, making up 90% of

the field, which can't be used for arable or livestock as the animals would damage the trees in their early years. David is hoping to sow it with wildflower seeds, as he has done elsewhere on the farm.

In 1998 a 35 acre arable field went into a 10 year ESA agreement as arable reversion to extensive grassland. The agreement was renewed in Autumn 2012 with added wildflowers from Cotswold Seeds. 'I like the website. It's very good and the staff are knowledgeable and helpful. The company is local so we can pick seeds up if we need them urgently, or there's 24 hour delivery.'

A contractor with years of experience of these types of mixes, sows with a grass seed drill using a low seed rate. The wildflower areas are later lightly grazed by the beef suckler herd. Management is low input, with no fertiliser. One year out of three it's cut for hay after the flowers have set seed. 'It's not grown as commercial grass and since the hay contains flowers it's not nutritionally as good as mainstream grasses,' David says. 'But the environmental benefits are really important for us. We've got beehives on the farm and they love the wildflowers. We have red kites and buzzards and barn owl boxes along the river banks. In May and June the grass is alive with insects.'

### **"It's a different way of looking at things, but for us commerciality needs to balance with environmental benefits"**

This is why David wants to increase the amount of wildflowers grown on the farm. Keeping the cricket bat willows as his primary cash crop, he is trialing equipment to make it more feasible to grow wildflowers around them. 'We hire contractors with big balers and machinery but we may have to downscale to manoeuvre between the trees. The flowers will stabilize soil while not competing with the trees.' Older grassy areas can be managed with Aberdeen Angus cattle which can use the coarse grass more efficiently.

David is also investigating growing wildflowers around the Christmas trees, which take 8 years to grow to a saleable height after planting. Cricket bat willows require even more patience, taking 15-18 years to mature. Blanks are selected and seasoned and sent to India.

'We're looking to use livestock to manage habitat as opposed to growing commercial grass to fatten livestock,' David explains. 'It's a different way of looking at things, but for us commerciality needs to balance with environmental benefits.'



# Wild Flowers

During the last decade, we've seen an increasing demand for wild flower seeds which are being sown to recreate traditional meadows which have been in decline.

Wild flower meadows are either managed under an agri-environmental agreement, where a list of species and management prescription will be provided by Natural England, or often for aesthetic purposes alone. They take many years to evolve naturally and can't be instantly created just by sowing seeds.

Nevertheless, with proper preparation and management, excellent results can be achieved in a relatively short time. **See our website for case studies and management advice.**

## Meadow Over-Seeding

Just Wild Flowers

Code: MIXWFOS

This wild flower-only mixture can be sown into open swards that are free of aggressive grasses and weeds. Sow in autumn when existing plant growth is slower.

- 12% lesser knapweed
- 10% native red clover
- 10% salad burnet
- 10% sainfoin
- 9% ox-eye daisy
- 7.50% ribwort plantain
- 7% white campion
- 6% musk mallow
- 5.50% self heal
- 5% betony
- 4.5% red campion
- 4% yarrow
- 3.50% wild carrot
- 3% ladys bedstraw
- 2% meadowsweet
- 0.50% small scabious
- 0.50% yellow rattle

2.00 kg/acre 5.00 kg/ha

£150.28 per kg +VAT



Cotswold wild flora  
Cheshire  
12th June

Image: Julian Kronfli Photography

## Mixes

### Cornfield Annuals

For One Summer

Code: MIXANN

This is a one year mix to provide a colourful display between June and August. Must be planted by April.

- 45% corn cockle
- 13% cornflower
- 13% corn marigold
- 14% field poppy
- 7% birdsfoot trefoil
- 4% crimson clover
- 4% alsike clover

6.00 kg/acre 15.00 kg/ha

£57.25 per kg +VAT

### Cotswold Wild Flora

Long Term

Code: MIXFLO

Our most popular mix combines annuals, for an exceptional display in the first year, with perennials which get better and better from year two onwards. Species included may vary occasionally.

- 5% certified common bentgrass
- 5% commercial sweet vernal grass
- 10% commercial crested dogstail
- 10% certified smaller catstail
- 15% certified red fescue
- 15% certified smooth stalked meadowgrass
- 25% certified sheeps fescue
- 1% salad burnet
- 1% lesser knapweed
- 1% self heal
- 1% ox-eye daisy
- 1% yarrow
- 1% betony
- 1% ladys bedstraw
- 0.5% ribwort plantain
- 0.5% meadowsweet
- 0.5% red campion
- 0.5% wild carrot
- 0.25% field scabious
- 0.25% small scabious
- 1.50% corn cockle
- 1% corn marigold
- 1% cornflower
- 1% field poppy
- 1% yellow rattle

10.00 kg/acre 25.00 kg/ha

£42.55 per kg



Grass

Wildflower

Legume

## Mixes

## Woodland Edge and Shady Area

## Long Term

Code: MIXWOOD

In open and semi-shaded areas a number of grasses and wild flowers will thrive many of which are in this mix.

- 1% quaking grass
- 1% commercial tufted hairgrass
- 2% commercial sweet vernal grass
- 8% certified common bentgrass
- 10% commercial crested dogtail
- 14% certified wood meadowgrass
- 24% certified slender creeping red fescue
- 25% certified red fescue
- 2.50% red campion
- 2.50% garlic mustard
- 1.50% self heal
- 1.50% tufted vetch
- 1% white campion
- 1% bluebell
- 1% yarrow
- 1% teasel
- 1% betony
- 0.75% wood avens
- 0.75% meadowsweet
- 0.25% autumn hawkbit
- 0.25% pignut

10.00 kg/acre 25.00 kg/ha

£48.45 per kg

## Acid &amp; Clay Soil

## Long Term

Code: MIXACID

A suitable mixture for both acidic and heavy clay soil types. Prepare a well worked, weed-free seedbed and spread seeds at no more than 10mm deep.

- 1% commercial quaking grass
- 4% commercial sweet vernal grass
- 10% commercial meadow foxtail
- 10% commercial crested dogtail
- 15% certified smaller catstail
- 20% certified common bentgrass
- 25% certified red fescue
- 2.5% lesser knapweed
- 2% betony
- 2% self heal
- 2% ox-eye daisy
- 1.25% common sorrel
- 1% ribwort plantain
- 1% lady's bedstraw
- 1% meadow buttercup
- 1% yarrow
- 0.5% native red clover
- 0.5% yellow rattle
- 0.25% sheep's sorrel

10.00 kg/acre 25.00 kg/ha

£54.36 per kg

## Chalk &amp; Limestone Soil

## Long Term

Code: MIXCHA

This mixture is designed for chalk and limestone soil types. Chalk and Limestone soils are known for their ability to support a large selection of wild flower species which is why we have been able to create such a diverse mixture.

- 5% certified common bentgrass
- 5% certified crested dogtail
- 10% certified smaller catstail
- 5% commercial yellow oatgrass
- 20% certified sheep's fescue
- 20% certified red fescue
- 20% certified smooth stalked meadowgrass
- 2.25% salad burnet
- 2% lesser knapweed
- 2% yellow carrot
- 1.50% meadow buttercup
- 1% lady's bedstraw
- 1% ox-eye daisy
- 1% ribwort plantain
- 1% self heal
- 1% yarrow
- 0.50% birdsfoot trefoil
- 0.50% agrimony
- 0.50% small scabious
- 0.50% field scabious
- 0.25% hoary plantain

10.00 kg/acre 25.00 kg/ha

£48.00 per kg

## Damp Meadow

## Long Term

Code: MIXDAM

Wetter soils require a slightly different seed mixture.

This one should give reliable results on most damp soils and may also be used around water courses or ponds.

- 2% commercial meadow foxtail
- 3.10% commercial crested dogtail
- 5% certified common bentgrass
- 6% certified smaller catstail
- 20% certified smooth stalked meadowgrass
- 25% certified red fescue
- 28% certified sheep's fescue
- 2.50% great burnet
- 1.50% ox-eye daisy
- 1.50% garlic mustard
- 1.20% devil's-bit scabious
- 1% lesser knapweed
- 1% common sorrel
- 1% self heal
- 0.50% yellow rattle
- 0.50% ragged robin
- 0.20% ribwort plantain

10.00 kg/acre 25.00 kg/ha

£52.60 per kg



# Wild Flower Directory

## Perennials

### Agrimony

*Agrimonia*

Upright plant found in hedges and field edges. Late seeding.



Flowers: June-Aug

Late

### Devil's Bit Scabious

*Succisa pratensis*

Found in damp meadows and wetter (but not waterlogged) areas.



Flowers: June-Sept

Late

### Meadow Buttercup

*Ranunculus acris*

Found in older grasslands and damp grassy places with a long flowering period.



Flowers: Apr-Oct

Early

### Betony

*Stachys officinalis*

Found in shady areas, woodland fringes & hedge rows. Likes damp sites.



Flowers: June-Sept

### Field Scabious

*Knautia arvensis*

Frequent in cornfields, grassland and roadsides on calcareous dry soils.



Flowers: June-Oct

Late

### Meadowsweet

*Filipendula ulmaria*

Found in and alongside meadows. Prefers wet ground. Strongly scented flowers.



Flowers: June-Sept

Late

### Birdsfoot Trefoil

*Lotus corniculatus*

Found in downlands and old pasture, esp. on calcareous soils, drought resistant.



Flowers: June-Sept

### Great Burnet

*Sanguisorba officinalis*

Oblong burgundy flower heads, found on wetter meadow ground.



Flowers: June-Sept

### Meadow Vetchling

*Lathyrus pratensis*

Yellow pea-like flower, grows in grassy fields and hedgerows.



Flowers: June-Sept

### Bluebell

*Hyacinthoides non-scripta*

Found in hedge-banks and woodland where they can form a distinctive blue carpet.



Flowers: Apr-June

Early

### Ladys/Hedge Bedstraw

*Galium verum/Galium mollugo*

Ladys bedstraw suits most soils. Hedge bedstraw prefers free-draining.



Flowers: June-Sept

### Musk Mallow

*Malva moschata*

Prolific on soils rich in nitrogen. Grows in hedgerows and grassland.



Flowers: June-Sept

### Cowslip

*Primula veris*

Found on chalky grassland and open calcareous woodland.



Flowers: Apr-May

Early

### Lesser Knapweed

*Centaurea nigra*

Also known as common or black knapweed. Good nectar source



Flowers: June-Sept

### Ox-Eye Daisy

*Leucanthemum vulgare*

Robust, reliable plant for alkaline soils. Found in meadows, pastures and banks.



Flowers: May-Sept



## Perennials continued

**Ragged Robin***Lychnis flos-cuculi*

Delicate ragged flowers usually found in damp meadows.

Early



Flowers: May-Aug

**Red Campion***Silene dioica*

Often found in woodland and shady areas. Likes damp soils.

Early



Flowers: May-Sept

**Ribwort Plantain***Plantago lanceolata*

Established in most older grassland. Source of vitamins and minerals for grazing animals.



Flowers: Apr-Oct

**St Johns Wort***Hypericum perforatum*

Likes free-draining calcareous soils with a sunny aspect. Has medicinal properties.



Flowers: June-Sept

**Salad Burnet***Sanguisorba minor*

Found on dry, lime rich, calcareous soils. Liked by grazing animals.



Flowers: May-Sept

**Self Heal***Prunella vulgaris*

A low growing, creeping plant which is common in most grassland.



Flowers: June-Oct

**Sorrel***Rumex acetosa*

Grows well in loamy soils rich in nutrients.



Flowers: May-Aug

**Teasel***Dipsacus fullonum*

A tall plant found in field margins, particularly in the south of Britain.



Flowers: July-Sept

**Tufted Vetch***Vicia cracca*

Creeping, sprawling growth habit. Found in hedgerows and climbing up vegetation.



Flowers: June-Aug

**White Campion***Silene latifolia*

Frequent in roadside verges, hedgerows and waste ground.



Flowers: May-Oct

**Wild Carrot***Daucus carota*

Found in grassy places, field margins and roadsides. Prefers calcareous soils.



Flowers: June-Sept

**Yarrow***Achillea millefolium*

Found in grassland and grass margins, hedgerows and open spaces.



Flowers: June-Nov

Late

## Annuals

**Corn Chamomile***Anthemis arvensis*

Corn field annual which thrives in loamy soils rich in nutrients.



Flowers: June-July

**Corn Cockle***Agrostemma githago*

A tall annual with an attractive vivid purple flower.



Flowers: May-Aug

**Cornflower***Centaurea cyanus*

A pretty bright blue solitary flower. Was used as a dye in champagne wine.



Flowers: June-Aug

**Corn Marigold***Crysanthemum segetum*

A former weed in spring-sown corn. Now rare on farmed land. Bold yellow flowers.



Flowers: June-Oct

**Field Poppy***Papaver rhoeas*

Found in arable fields and disturbed ground. Silky, deep scarlet flowers.



Flowers: June-Oct

**Yellow Rattle***Rhinanthus minor*

Parasitic plant which restricts grass growth allowing delicate wildflowers to establish.



Flowers: May-Sept

\*Please note wild flower seed sold as straight attracts VAT at the current rate of 20%





# Game

Reliable game cover and food for any shoot.

The game cover section has been recently updated, it provides a wider choice of mixtures that have been in development for the last 3 years. The FlexiCover mixtures provide both 1 and 2 year options and can be used for both flushing and holding cover. The combination of grain sorghum and brassica species provides reliable cover all the way through the winter.

The new Cotswold Partridge mix combines species attractive to partridge and a broken canopy to protect against predators from above. While our range is more comprehensive than ever before, no one mixture will fit all shoots and sites, so we are more than happy to talk through different species and options and tailor bespoke mixtures to fit.

## Game and Bird Food Crop Overview

Species	Duration	Sowing Time	Sowing Depth	Full Growth Height (cm)	Comments	Sowing Rate (kg/ac)	Feed	Cover
Maize	1 Yr	Late April-May	7.5 - 10	180 - 200	Cobless varieties available	11 - 14	✓	✓
R. Millet	1 Yr	April-June	2.5	100 - 120	Later seeding than W.Millet	5 - 10	✓	
W. Millet	1 Yr	April-June	2.5	100 - 120	Produces more seed than R. Millet	5 - 10	✓	
Reed Millet	1 Yr	April-June	2.5	100 - 120	Strong standing ability	5 - 10		✓
D. Sorghum	1 Yr	May-June	3 - 5	100	Sow in wide rows	8		✓
Giant Sorghum	1 Yr	May-June	3 - 5	180	Prone to brackling	12		✓
Grain Sorghum	1 Yr	May-June	3 - 5	100 - 120	Produces seed	8	✓	✓
Sunflower	1 Yr	Mid April onwards	5	90 - 175	Dwarf varieties reach 3ft	10	✓	
Buckwheat	1 Yr	May-June	3.5	90	Not frost hardy	20 - 30	✓	✓
Linseed	1 Yr	March-June	2	50 - 60	Good for Partridge	20	✓	✓
S. Cereals	1 Yr	March-May	2 - 3	70 - 80	Sow in spring for winter grain	50 - 75	✓	✓
W. Cereals	1 Yr	March-Sept	2 - 4	70 - 90	Sow in autumn for grain in Yr 2	50 - 75	✓	✓
Quinoa	1 Yr	May-June	0.5 - 1	90 - 140	Produces high protein seed	5	✓	
F. Rape/OSR	1 Yr	May-August	1	80 - 90	Flea beetle risk	4		✓
Mustard	1 Yr	May-August	1	80 - 120	Sow in august for late cover	6 - 10	✓	✓
Brown Mustard	1 Yr	May-August	1	80 - 100	More winter hardy than Mustard	2	✓	✓
Fodder Radish	1 Yr	May-August	1	80 - 120	Holds seeds late in season	6	✓	✓
Hybrid Brassica	1 Yr	April-August	1	90 - 120	Sow by mid Aug	3		✓
Gold of Pleasure	1 Yr	April-May	1	50 - 70	High seed shed	5	✓	✓
Kale	2+ Yr	April-June	1	70 - 110	2 year cover	3		✓
Sweet Clover	2+ Yr	April-June	0.5 - 1	120	Significant growth in Yr 2	6		✓
Chicory	2+ Yr	April-Sept	0.5 - 1	90 - 150	Lasts 3-4 Years	6	✓	✓
Canary Grass	2+ Yr	May-June	1	180	Main growth in Yr 2 onwards	3	✓	
Reed C. Grass	2+ Yr	May-June	1	200	More winter hardy than Canary Grass	3	✓	



## Mixes

## FlexiCover One Year Game Mix

## Cover and Feed

Code: MIXFLEXI

This flexible mixture combines brassicas, sorghums and cereals. Sowing in wide rows allows game birds easier movement if pushing them into a flushing point or sow in narrow rows to create a denser holding cover, or windbreak alongside maize.

- 3.85 kg certified grain sorghum
- 3.80 kg certified spring triticale
- 3.50 kg certified spring barley
- 1.10 kg red millet
- 1.10 kg white millet
- 0.40 kg reed millet (Japanese)
- 0.70 kg certified forage rape
- 0.60 kg certified forage rape/kale hybrid
- 0.50 kg certified fodder radish
- 0.20 kg commercial gold of pleasure
- 0.20 kg certified mustard
- 0.05 kg certified brassica carinata

16.00 kg/acre - £46.99

40.00 kg/ha - £117.48

## FlexiCover Two Year Game Mix

## Cover and Feed

Code: MIXFLEX2

The inclusion of kale can ensure this mixture lasts for two full years. During establishment protect against flea beetle and consider fertiliser to push the brassicas past the most susceptible stage of pest damage.

- 4.00 kg certified spring triticale
- 3.00 kg certified spring barley
- 3.25 kg certified grain sorghum
- 1.10 kg red millet
- 1.10 kg white millet
- 0.30 kg reed millet (Japanese)
- 2.15 kg certified game kale
- 0.40 kg certified fodder radish
- 0.20 kg certified forage rape/kale hybrid
- 0.10 kg commercial gold of pleasure
- 0.40 kg fennel

16.00 kg/acre - £63.15

40.00 kg/ha - £157.88

## Retrieve Mix

## Fast and Economical

Code: MIXRET

For a summer sowing after a failed spring crop nothing beats these fast growing brassica species. It's quick, reliable and it works.

- 2.88 kg mustard
- 1.56 kg forage rape
- 0.96 kg fodder radish
- 0.48 kg hybrid rape/kale
- 0.12 kg brassica carinata

6.00 kg/acre - £19.90

15.00 kg/ha - £47.75

## Mixes

## General Purpose Game Mix

## Cover and Feed

Code: MIXGAME

This is our most popular game cover mix, combining a wide range of species providing feed and cover for pheasants, partridge and farmland birds.

- 2.25 kg commercial buckwheat
- 2.00 kg commercial sunflower
- 1.30 kg white millet
- 1.30 kg red millet
- 0.50 kg certified forage rape
- 0.50 kg certified mustard
- 0.50 kg reed millet (Japanese)
- 0.50 kg certified fodder radish
- 0.30 kg quinoa
- 0.25 kg certified hybrid rape/kale
- 0.30 kg certified game kale

10.00 kg/acre - £39.90

25.00 kg/ha - £99.75

## Cotswold Partridge Mix

## Cover and Feed

Code: MIXPART

The mix is designed to include species which attract partridge, as well as creating a broken canopy with room for birds to move through the cover.

- 6.50 kg certified spring triticale
- 3.60 kg certified spring wheat
- 3.60 kg certified spring barley
- 1.45 kg white millet
- 1.45 kg red millet
- 1.45 kg certified linseed
- 0.45 kg certified forage rape
- 0.45 kg certified gold of pleasure
- 0.20 kg certified hybrid rape/kale
- 0.20 kg certified leafy turnips
- 0.20 kg certified crimson clover
- 0.45 kg fennel

20.00 kg/acre - £43.79

50.00 kg/ha - £109.48



Seed and shelter millet mix  
FarmED  
10th September



Mixes

Quinoa/Kale Mix

Cover and Feed Code: MIXQUI

This simple combination supplies the two key requirements of birds: cover and feed. The kale provides excellent winter cover and supports the quinoa plants. Quinoa can provide 1-2t per acre of high protein feed from late autumn.

- 1.40 kg quinoa
- 1.45 kg game kale
- 0.15 kg fodder rape/kale hybrid

3.00 kg/acre - £35.35 7.50 kg/ha - £88.38

Short & Sturdy Game Cover Mix

Cover and Feed Code: MIXSHORT

This mix is ideal for growing with maize to act as a windbreak on exposed sites or to create a flushing point in front of the gun line. The seed bearing grain sorghum, sunflower and millet also provide for farmland birds.

- 4.70 kg commercial grain sorghum
- 2.50 kg certified dwarf sunflower
- 0.50 kg white millet
- 0.30 kg reed millet (Japanese)

8.00 kg/acre - £36.58 20.00 kg/ha - £91.45

Seed & Shelter Millet Mix

Cover and Feed Code: MIXMIL

Combining Red and White Millet for a wider window of seed production and reed millet to provide cover. Broadcast or shallow drill in maize strips for a denser cover.

- 3.20 kg white millet
- 2.80 kg red millet
- 2.00 kg reed millet (Japanese)

8.00 kg/acre - £28.10 20.00 kg/ha - £70.25

Sowing and Growing

Whether you run a small local syndicate or a large estate shoot it is important to produce good, reliable crops that provide plenty of shelter, cover and seed.

When to sow

Most game crops are spring sown after frost risk has passed to provide cover and feed from late summer. Start planting the mixes of brassicas, millet, maize, sunflowers and canary grass in mid April, with dwarf sorghum better if drilled in May or June. The only exception is the quick-growing Retrieve Mixture which can be drilled anytime from April to early September if there is sufficient soil moisture.

How to sow

A well worked weed-free seedbed is required, try to achieve a stale seed bed with several cultivation passes to stimulate weed germination before sowing. Seeds such as maize, sunflower and sorghum are usually drilled but small seeded species such as kale and mustard may be broadcast and well rolled after sowing.

If sowing a mix with a range of seed sizes, prioritise the smaller seeds, try to ensure the smaller seeds are not sown too deeply, as this can reduce the reliability of establishment, most larger seeds will cope with being sown slightly shallower.

Bespoke Autumn sown mixtures are available for cover in the following year.

Management

For sites with a known weed burden of later germinating annuals like fat hen consider herbicide tolerant mixtures (Speak to an adviser). Grass weeds can also be controlled by sowing broadleaf only mixtures tolerant to graminicides.

Nutrient requirements

Game crops require P & K levels to be ADAS Index 2 and benefit from 50-100kg N/ha in the seed bed. Farmyard manure can also be a very beneficial fertility source which will break down over several years.



## Straights

## Game Maize

Specifically chosen for its early establishment vigour and impressive standing ability, providing robust cover late in the season. This variety will reach 2 metres and has medium to late maturity. If the priority is early cob production over holding cover please contact us for further options. **Only available in one acre packs (treated seed).**

**Certified Game Maize     £42.50 per acre**

## Cotswold Game Kale Blend

The kale mix combines the tall and leafy Proteor variety with the shorter, strong Keeper kale variety, creating shelter with space for birds to move through the cover.

**3.00 kg/acre - £34.95     7.50 kg/ha - £87.38**

## Dwarf Sorghum

An excellent windbreak around other crops, it is an annual crop with similar properties to maize. **Only available in one acre packs.**

**Certified Dwarf Sorghum     £36.80 per acre**

## Straights

## Canary Grass

This perennial grass is drilled in wide rows (60-90cm) and takes a year or so to become established. From the second year the seed heads will reach two metres and the crop can usually be relied upon for 10 years. Good for pheasants and partridges as well as linnets and wrens.

**3.00 kg/acre - £54.30     7.50 kg/ha - £135.75**

## Reed Canary Grass


Grows taller than Canary grass and is more suitable to northern climates because it tolerates extreme cold weather. Reed canary grass can be slow to establish, sometimes taking up to two years but it will tolerate a wide range of soil types.

**3.00 kg/acre - £88.05     7.50 kg/ha - £220.13**

## Sunflower

Large amounts of food supplied through the winter. Please specify dwarf (1-2 metres) or standard type (2 metres+).

**10.00 kg/acre - £47.50     25.00 kg/ha - £118.75**







# Lawn & Landscape

Seed mixtures that establish quickly to provide attractive, tough and low maintenance turf.

Great lawns are a distinctive feature of British gardens. Our mixtures contain the best modern varieties that require minimal maintenance so you can enjoy more time looking and less time mowing.

The right mix is just as important for other key grassy areas such as sports pitches and roadside verges. Creating the right species mix for these uses is vital to ensure the surface is correct for purpose and can be effectively managed.

## How Much Seed?

- ▶ Measure the area of your lawn in square metres (multiplying the width by the length will give the area).
- ▶ Multiply the area by the sowing rate given for each mix (in grams per square metre), always using the highest sowing rate if you are creating a lawn on bare earth
- ▶ Divide the results by 1000 to give the number of kilograms required

Example:

Length of lawn = 10m, width = 4m, area of lawn = 40m<sup>2</sup>  
 40 x 70g (sowing rate) = 2800 = 2.8kg of seed required

Please note: One acre = 4000m<sup>2</sup>

	70 grams/m <sup>2</sup> High rate	50 grams/m <sup>2</sup> Medium rate	35 grams/m <sup>2</sup> Low rate
15 m <sup>2</sup>	1 kg	0.75 kg	0.5 kg
30 m <sup>2</sup>	2 kg	1.5 kg	1 kg
100 m <sup>2</sup>	7 kg	5 kg	3.5 kg
250 m <sup>2</sup>	17.5kg	12.5 kg	8.75 kg
500 m <sup>2</sup>	35 kg	25 kg	17.5 kg

## Mixes: Lawn

### Hard Wearing Lawn

With Ryegrass

Code: MIXHAR

Designed to produce a tough and durable lawn which is easy to grow and live with. The grasses used produce a knitted turf and offer unbeatable toughness. We have supplied this mixture for 30 years and frequently hear positive reports from our customers. As well as being used as a lawn, this mixture will produce a suitable turf for caravan parks and airfields. Turf growers also like the blend as it establishes quickly.

- 50% certified dwarf/turf ryegrass
- 40% certified slender creeping red fescue
- 10% certified common bentgrass

50 - 70 g/m<sup>2</sup>

£5.96 per kg

### Ornamental Lawn

Without Ryegrass

Code: MIXORN

A superb fine-leaved mixture with a superior finish. It is suitable for sites where a 'bowling green' finish is required. The mixture is slow growing and tolerates light shade but will require regular feeding and mowing, ideally with a cylinder mower, to keep its appearance.

- 80% certified slender creeping red fescue
- 20% certified common bentgrass

50 - 70 g/m<sup>2</sup>

£6.75 per kg

### Shady Lawn

Without Ryegrass

Code: MIXSHA

A slow-growing mixture for low-light areas. A good turf can be produced from this mix, especially when the mowing height is increased to 2-3 inches.

- 30% certified slender creeping red fescue
- 30% certified red fescue
- 30% certified smooth stalked meadowgrass
- 10% certified common bentgrass

50 - 70 g/m<sup>2</sup>

£6.20 per kg



## Mixes: Landscape

## Economy Landscape

## With Ryegrass

Code: MIXECO

A low cost, fast establishing mixture, designed to cover large areas quickly. Suitable for banks and other non-prestigious areas.

- 50% certified dwarf/turf ryegrass
- 50% certified creeping red fescue

35 g/m<sup>2</sup>

£4.68 per kg

## Verge Mixture

## With Ryegrass

Code: MIXRV

An all-round mixture which is used for verges, pipelines and other reinstatement projects.

- 35% certified dwarf/turf ryegrass
- 30% certified creeping red fescue
- 20% certified chewings/red fescue
- 10% certified smooth stalked meadowgrass
- 5% certified common bentgrass

35 - 70 g/m<sup>2</sup>

£5.11 per kg

## Low Maintenance

## Without Ryegrass

Code: MIXLM

A seed mixture which requires very little mowing or maintenance. It will tolerate shade and grows on all soils including infertile types.

- 70% certified sheeps fescue
- 20% certified chewings/red fescue
- 10% certified common bentgrass

35 - 70 g/m<sup>2</sup>

£5.47 per kg

## Additions



## White Clover

Nitrogen fixing clover.

White clover can be included on request.

**Please call for advice.**

## Mixes: Sport

## Rugby &amp; Football

## Re-Seed and Renovate

Code: MIXSPO

This is a fast establishing mixture which can be used for all winter sports. It is also suitable for school playing fields and for over-seeding pitches at the end of the season.

- 80% certified dwarf/turf ryegrass
- 20% certified creeping red fescue

35 - 50 g/m<sup>2</sup>

£4.58 per kg

## Tennis Court &amp; Cricket Wicket

## With Ryegrass

Code: MIXTEN

Ball bounce and speed are improved on dense and compact turf. Formulated to provide a firm and fast surface and give a true, sufficiently high bounce. Frequent mowing and rolling recommended for best results.

- 45% certified dwarf/turf ryegrass
- 30% certified chewings/red fescue
- 15% certified slender creeping red fescue
- 10% certified common bentgrass

50 g/m<sup>2</sup>

£5.64 per kg



Hard wearing lawn  
1st June





Available to request or download at [cotswoldseeds.com](https://cotswoldseeds.com)

## COTSWOLD SEEDS

Cotswold Seeds was founded in 1974 and deals with over 17,000 farmers throughout the UK. The company has a specialist interest in grass and legumes, offering advice on growing and managing these crops to farmers and growers in the livestock, arable and horticultural sectors. The company, in conjunction with FarmED at Honeydale Farm, is also involved in a wide range of research projects.

Cotswold Seeds Ltd  
Cotswold Business Village  
Moreton in Marsh  
Gloucestershire, GL56 0JQ

Tel: 01608 652552  
[www.cotswoldseeds.com](https://www.cotswoldseeds.com)  
[info@cotswoldseeds.com](mailto:info@cotswoldseeds.com)

ISBN 978-1-8380163-1-9  
DEFRA Reg. No. 165  
BSPB Licence No. 1476  
Soil Association Licence No. P5985  
Registered in England 1163604  
VAT No. 195 5792 09

Follow us on:



[twitter.com/cotswoldseeds](https://twitter.com/cotswoldseeds)  
[facebook.com/cotswoldseeds](https://facebook.com/cotswoldseeds)  
[youtube.com/cotswoldseeds](https://youtube.com/cotswoldseeds)  
[instagram.com/cotswoldseeds](https://instagram.com/cotswoldseeds)

